This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

3 2044 106 371 867

LIBRARY OF THE GRAY HERBARIUM
HARVARD UNIVERSITY
THE GIFT OF

OCCASIONEM COGNOSCE.

John Amory Lowell.

Per
ENG
E-1

HARVARD UNIVERSITY HERBARIUM.

THE GIFT OF

John Amory Lowell.

MDCCCLXV.

EDWARDS'S

BOTANICAL REGISTER:

OR,

ORNAMENTAL FLOWER-GARDEN AND SHRUBBERY:

CONSISTING OF

COLOURED FIGURES OF PLANTS AND SHRUBS,

CULTIVATED IN BRITISH GARDENS;

ACCOMPANIED BY THEIR

History, Best Method of Treatment in Cultivation, Propagation, &c.

◆

CONTINUED

BY JOHN LINDLEY, Ph. D. F.R.S. L.S. AND G.S.

PROFESSOR OF BOTANY IN THE UNIVERSITY OF LONDON,
AND THE ROYAL INSTITUTION OF GREAT BRITAIN.
&c. &c. &c.

New Series. Pl. 1742-1821, 1835
1822-1828, 1836

VOL. VIII. cf. Journ. Arn. Arb. 18:
184. 1937.

OR VOL. XXI. OF THE ENTIRE WORK.

——viret semper——nec fronde caducâ
Carpitur.

LONDON:
JAMES RIDGWAY AND SONS, PICCADILLY.

M.DCCC.XXXVI.

ALPHABETICAL INDEX

TO

VOL. VIII. OF THE NEW SERIES.

Folium

Angræcum distichum 1781
Angræcum micranthum 1772
Arctostaphylos tomentosa 1791
Arbutus procera 1753
Aristolochia fœtens................ 1824
Azara dentata 1788
Begonia petalodes 1757
Berberis dealbata 1750
Bletia reflexa 1760
Brassia Lanceana 1754
Calceolaria angustiflora 1743
Calotropis procera 1792
Campanula garganica 1768
Caprifolium hispidulum 1761
Cereus triangularis 1807
Chironia peduncularis 1803
Clavija ornata 1764
Clianthus puniceus 1775
Coccoloba virens 1816
Colletia horrida 1776
Coryanthes maculata 1793
Cosmelia rubra 1822
Cratægus Douglasii 1810
Cycnoches Loddigesii 1742
Daubenya aurea 1813
Dendrobium Pierardi 1756
Dendrobium cupreum 1779
Dendrobium densiflorum 1828
Dracæna terminalis 1749
Dyckia rariflora 1782
Edwardsia chilensis 1798
Elichrysum bicolor 1814
Empetrum rubrum 1783
Epidendrum gracile 1765
Eriogonum compositum 1774
Erythronium grandiflorum 1786
Eutoca divaricata 1784
Eutoca viscida 1808
Fernandezia acuta 1806
Fuchsia discolor 1805
Galatella punctata 1818
Gardoquia Gilliesii 1812
Gardoquia Hookeri 1747

Folium

Gesneria allagophylla 1737
Gesneria faucialis 1785
Govenia superba 1795
Hibiscus Rosa sinensis 1826
Indigofera atropurpurea 1744
Ipomœa Aitoni 1794
Kennedya Marryattæ 1790
Lælia anceps 1751
Lasthenia californica 1823
Lasthenia glabrata 1780
Lepanthes tridentata 1762
Macradenia triandra 1815
Maxillaria cristata 1811
Maxillaria crocea 1799
Maxillaria densa 1804
Maxillaria picta 1802
Mimulus luteus, var. *variegatus* 1796
Monachanthus viridis 1752
Myanthus barbatus 1778
Ochranthe arguta 1819
Oncidium citrinum 1758
Oncidium Lemonianum 1789
Oncidium pulchellum.............. 1787
Orobus atropurpureus 1763
Oxalis Piottæ 1817
Pæonia Moutan lacera 1771
Pentstemon staticifolius 1770
Pholidota imbricata 1777
Physianthus albens 1759
Pimelia ligustrina 1827
Pleurothallis Grobyi 1797
Pleurothallis picta 1825
Prunus japonica 1801
Psoralea macrostachya 1769
Rhodochiton volubile 1755
Rhododendron pulcherrimum....... 1820
Russellia juncea 1773
Stanhopea oculata 1800
Tritoma Burchelli 1745
Vanda teres 1809
Verbena multifida contracta........ 1766
Verbena sulphurea 1748
Zephyranthes Spofforthiana 1746

1742.

1742.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly March 1. 1835. S. Watts sc.

1742

*CYCNÓCHES Loddigèsii.

Loddiges' Swanwort.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § VANDEÆ, (*Introduction to the Natural System of Botany*, p. 262.)

CYCNOCHES, Lindl. Perianthium explanatum. *Sepala* lateralia lanceolata, basi paululùm sub labello connata; supremo angustiore. *Petala* latiora, falcata, decurva. *Labellum* liberum, ecalcaratum, columnâ continuum, lanceolatum, integerrimum, ungue abrupto calloso. *Columna* elongata, arcuata, teres, apice clavata, auriculis 2 falcatis ad latera clinandrii. *Anthera* bilocularis. *Pollinia* 2, postice sulcata, subpedicellata; *caudiculâ* lineari; *glandulâ* grossâ.—Habitus *Cataseti* (sed racemus lateralis). *Lindl. Gen. & Sp. Orch.* p. 154.

Cycnoches Loddigesii, *Lindl. l. c. Bot. Cab. t.* 2000.

Caulis *erectus, carnosus, cylindraceus, subpedalis, pallidè viridis, corrugatus, basi vaginis membranaceis, apice foliis vestitus.* Folia *omninò* Cataseti *cujusdam.* Racemus *multiflorus, pendulus, e latere caulis erumpens, flexuosus.* Flores *maximi ordinis, initio inodori, mox Vanillam gratissimè spirantes, præsertim manè.* Bracteæ *ovatæ, cucullatæ, obtusæ, pedicellis breviores.* Sepala *viridia apice fusca, et obscure maculata, mox rufescentes; supremum lineari-oblongum, arcuatum 3-pollicare; lateralia 2-poll. longa imâ basi connata, ovato-oblonga, subundulata, pendula.* Petala *ejusdem coloris cum sepalis, sed minùs conspicuè maculata, oblongo-lanceolata, inæquilatera, decurva, 2½-poll. longa.* Labellum *patentissimum, cum columnâ continuum, angustè oblongum, convexum, carnosum, medio album, apicem versus tenuius, sordidè lutescens, undique maculis latis inæqualibus sanguineis notatum;* ungue *brevi alato maculoso, alis lutescentibus, disco convexo albo.* Columna *gracillima, arcuata, utrinque clavata; dimidio inferiore atropurpureo, superiore complanato virescente purpureo maculoso.* Anthera *citò decidua, membranacea, bilocularis, ferè pellucida, omninò mutica.* Clinandrium *posticè bicorne, cornubus falcatis compressis supra antheram incurvis.*

For the discovery of this most extraordinary Epiphyte the public is indebted to John Henry Lance, Esq. who discovered it, and a great number of other new plants in the woods

---

* So called from κύκνος a swan, and αὐχὴν the neck; in allusion to the column of this plant, which is gracefully curved, like the neck of the swan.

of Surinam. It was originally sent by that gentleman to Messrs. Loddiges, with whom it flowered imperfectly three or four years ago. Other and more healthy plants were afterwards presented by Mr. Lance to the Horticultural Society, the produce of one of which was the singular raceme now represented.

In size the flowers of this plant are the largest that have yet been met with among Orchideous plants, measuring not less than five inches from the tip of the back sepal, to the point of the lip. They are more singular than beautiful, but they compensate in fragrance for their want of brilliant colours: the purest odour of Vanilla is exhaled by them when they have been open a short time, especially in the morning.

This species flowers in the months of June and July. It resembles a Catasetum in habit, but its flowers originate from the side, and not the base, of the fleshy stem. It grows very freely under the same management as Catasetums, but seems to require the greatest heat that is ever obtained in a damp stove. It is disposed to increase itself pretty freely, by multiplication of its fleshy stems, and will probably cease, after a few years, to be so rare a plant as it now is.

1743

1743



rake. del. Pub by J. Ridgway 169 Piccadilly March 1 1835. J. Watts sc.

* CALCEOLÁRIA angustiflòra.

Narrow-flowered Calceolaria.

DIANDRIA *MONOGYNIA.*

Nat. ord. SCROPHULARINEÆ *Juss.* (*Introduction to the Natural System of Botany, p.* 228.)

CALCEOLARIA.—Supra, vol. 9. *p.* 723.

C. *angustiflora;* caule perenni diffuso, foliis cordato-ovatis duplicato-serratis glabris oppositis ternatisque, pedunculis axillaribus paucifloris sæpius paniculam angustissimam foliosam efficientibus, corollæ labio inferiore obtuso basi angustato superiore minimo.

C. angustiflora. *Ruiz et Pavon. Fl. Peruv. vol.* 1. *p.* 17. *t.* 28. *f. a. Graham in Bot. Mag. t.* 3094.

? C. verticillata. *Hooker Bot. Misc. vol.* 2. 233.

A half hardy perennial, flowering in August and September, and remarkable among its race for its prostrate habit, deep green, somewhat shining leaves, and small flowers. It was found first by the authors of the Flora Peruviana, in the valley of Canta in Peru; subsequently Mr. Cruckshanks discovered it at Culluay in the same district, and gathered the seeds from which the plants in our gardens were obtained.

It is a species of no great attraction, but deserves to be recorded in this work, as one of the genuine wild forms of a genus, which, however beautiful and interesting, has already begun to sink in estimation, in consequence of the ruin that has been brought upon it by the unskilfulness of gardeners. In their haste to improve the works of nature, these gentlemen have converted some of the fairest races in the Vegetable world, into forms in no case more beautiful than the original, and in the majority of instances unhealthy, mongrel, and debased. We strongly recommend all those who value this really beautiful and most singular genus, to abandon a pursuit which has as yet led to few results of which good taste can approve, and to apply the same skill which they have used in spoiling Calceolarias to recovering the pure original races, to preserving them uncontaminated, and to increasing their native charms ; not by unnatural combinations, but by those well known methods by which the purity of a species may be maintained while its vigour, health and beauty are augmented.

---

* See fol. 1214.

At least, if the genus must be the subject of hybridizing, let the intermixture be made with some reasonable attention to the only rules by which it is possible to arrive at a really desirable result.

1744

1744



Miss Drake del.   Pub by J. Ridgway 169 Piccadilly March 1. 1835.   J. Watts sc.

*INDIGÓFERA atropurpúrea.

Purple-flowered Indigo plant.

DIADELPHIA *DECANDRIA.*

Nat. ord. LEGUMINOSÆ *Juss. (Introduction to the Natural System of Botany, p.* 86.)

INDIGOFERA.—Supra, vol. 5. *fol.* 386.

I. *atropurpurea;* fruticosa, erecta, foliis 5-6-jugis, foliolis oblongis obtusissimis apiculatis tenuibus glabris, racemis multifloris foliis æqualibus v. longioribus, leguminibus linearibus rectis acustriatis 8-9-spermis.

I. atropurpurea *Hamilt. in Roxb. Fl. Ind. vol.* 3. 380. *Wall. Cat. No.* 5463.

A native of Nepal, where it was originally found by Hamilton; probably in the hot valleys; for it does not succeed well, unless cultivated in the stove. There it becomes a handsome light green bush, richly ornamented by its numerous bunches of purple and crimson flowers. If planted in the open air, it languishes even on a south wall and in such a season as that of 1834; producing its blossoms in small quantities and imperfectly.

Our specimens were communicated in August, 1833, from the hothouse of James Bateman, Esq. of Knypersley.

Roxburgh says, that when raised in the Botanic Garden at Calcutta, it grew from 3 to 5 feet high in about 8 months.

---

* Literally *Indigo-bearer;* some species of this genus producing the dye of that name.

174

1745.



Miss Drake del.　Pub by Ridgway 169 Piccadilly March 1.1835.　S. Watts sc.

1745

*TRÍTOMA Burchéllii.

Burchell's Tritoma.

HEXANDRIA *MONOGYNIA.*

Nat. ord. Asphodeleæ *Juss.* (*Introduction to the Natural System of Botany, p.* 273.)

TRITOMA Ker. Link. *Perianthium* tubulosum, eplicatum; limbo brevi 6-dentato regulari. *Stamina* hypogyna, recta, libera, exserta, alterna longiora. *Stigma* simplex. *Capsula* cartilaginea, rigescens, ovata, obtusè trigona. *Semina* biserialia, plurima, triquetra v. angulata, sibi invicem appressa.—Folia *graminea canaliculata.* Caulis *nullus.*

T. *Burchellii;* foliis lætè viridibus margine lævibus, racemo oblongo crasso denso, perianthiis clavato-cylindraceis unicoloribus.

Tritoma Burchellii. *Herbert. in Sweet's Hort. Brit.*

Racemus 4 *poll. longus, densissimus, et vix angustior, nullo modo pyramidatus v. protractus.* Flores *juniores sanguinei, posteà lutei v. potius luteo-aurantiaci unicolores,* 1¼-*poll. longi; limbo* 4 *lin. tubo apice* 3 *lin. lato.*

Communicated by the Honourable and Rev. William Herbert in July last. It was originally introduced from the Cape of Good Hope by Mr. Burchell, after whom it has been named.

It is a beautiful perennial plant, and quite hardy. At Spofforth it always flowers before or soon after Midsummer; all the other species of Tritoma are late flowerers.

Increased by the offsets, which it throws up in plenty.

---

* So named from τρεις three, and τεμνω to cut, because of the leaves ending in three sharp angles. As there was a genus of insects already called Tritoma, when this genus of plants was first proposed, Professor Link would change its name to Tritomanthe; but there is no end to alterations of this sort.

1746

1746.



[illegible] Pub. by S. Curtis [illegible] Walworth Mar. 1 1815. J. Watts sc.

* ZEPHYRÁNTHES Spofforthiána.

Spofforth Zephyranthes.

HEXANDRIA *MONOGYNIA.*

Nat. ord. AMARYLLIDEÆ, *R. Br.* (*Introduction to the Natural System of Botany, p.* 259.).

GARDEN VARIETY.

ZEPHYRANTHES SPOFFORTHIANA. Herbert MSS.

"Scape yellowish green, 5¾ inches high. Spathe brownish green. Peduncles green, an inch long. Germ deep green. Tube green. Limb rose-coloured, lined with white in the form of a star. Style very much declined, white and much longer than the filaments. Leaves not ¼ of an inch wide, acute, keeled, deep green. Pollen abundant, and anthers quite perfect and apparently fertile."

"The pretty flower which is the subject of this article, is an hybrid production from the tropical *Z. tubispatha*, which has white flowers, fertilized by the pollen of the Mexican *Z. carinata* with large red flowers. It is in every respect intermediate, having the leaves wider than those of *Z. tubispatha,* and distinguishable by the keel, which is conspicuous on the back of those of the male parent. The flower is also of intermediate dimensions and colour, resembling the male parent most both in form and hue, and in the posture of the style and filaments. A single bulb of this intermixture was obtained at Spofforth, which has produced a few offsets, piercing the ground, like those of the male parent, at a little distance from the principal bulb, their lateral direction giving the plant a stoloniferous appearance. Its anthers are perfectly formed, and the pollen seems to have all the requisites for fertility. Standing in a cool part of the stove, beside *Z. tubispatha*, it flowers before it, and probably will succeed like *Z. carinata* in a lower temperature. It must be observed that *Z. carinata*, Bot. Mag. the male parent of this plant, is, with respect to the flower only, the plant represented in the Bot. Reg. 902. under the name of *Z. grandiflora*, though the flower in the figure is rather longer and paler than its usual habit; but in consequence of the flower having been produced by a a newly imported bulb, before the leaves had shot, and a subsequent mistake of the gardeners, the sketch of the leaves, and the description of the leaves and seeds of *Z. grandiflora* were made erroneously from a pinkish variety of the small flowering *Z. striata*.

---

* See folio 1361.

"It will have been observed that *Habranthus phycelloides* of this work approaches so near in its general aspect to the genus *Phycella*, as to make it questionable whether the genus *Phycella* should be maintained distinct from *Habranthus*, at least from the many-flowered, non-expanding portion of that genus; and that the one-flowered *Habranthi*, which expand in the sun, are not easily distinguishable from the genus *Zephyranthes*; and it may be found that such species, viz. *H. versicolor*, *robustus*, and *Andersoni*, should be removed to the genus *Zephyranthes*; but an accurate revision of the various species should precede any alteration. It has, however, been as yet found impossible to obtain a mule between any *Habranthus* and *Zephyranthes*, which gives some reason to believe that the division as it stands is correct. Several flowers of *Z. striata* at Spofforth have lately been fertilized by *Z. carinata*, and others in the same pots by *H. Andersoni*, all the former appear to be forming seed, and all the latter have died away."

For the foregoing interesting memorandum we are indebted to the Honourable and Rev. William Herbert. We have for some time been aware of the error to which he alludes, and are happy to have the present opportunity of making it known. It will be necessary to expunge the name of *Z. grandiflora* from the catalogues, as it is a nonentity.

1747

1747.



[illegible] Piccadilly March 1 1815 J. Watts sc.

1747

* GARDOQUÍA Hookéri.

Scarlet Gardoquia.

DIDYNAMIA *GYMNOSPERMIA.*

Nat. ord. LABIATÆ, *Juss.* (*Introduction to the Natural System of Botany*, *p.* 239.)

GARDOQUIA Fl. Per. Calyx tubulosus, 13-nervius, subincurvus, ore æquali vel obliquo, dentibus brevibus rectis subæqualibus vel subbilabiatis. *Corollæ tubus* longè exsertus, rectus vel incurvus, intùs nudus; limbus bilabiatus, labium superius erectum subplanum emarginatum, inferius subpatens, lobis planis medio latiore. *Stamina* 4, subdidynama, inferioribus longioribus, laxè adscendentia, apice subdistantia, superiora subinde sterilia. *Filamenta* edentula. *Antheræ* biloculares, loculis distinctis parallelis vel subdivergentibus. *Styli* lobi subæquales. *Achenia* sicca, lævia.——Suffrutices fruticesve *ramosissimi, foliosi, sæpe procumbentes.* Flores *pulchri, sæpiùs coccinei.* Bentham Gen. et Sp. Lab. p. 397.

G. *Hookeri*; fruticosa, glaberrima, foliis obovatis subintegerrimis basi in petiolum brevem angustatis utrinque viridibus, pedunculis 2-3-floris, calycis glabri subbilabiati labio superiore breve bidentato erecto, fauce intùs villis clausâ. *Bentham l. c.* 401.
Cunila coccinea, *Nutt. ex Hooker exot. fl.* 3. *t.* 163.
Melissa coccinea, *Spreng. syst.* 2. 224.
Gardoquia Hookeri, *Don in Sweet. Fl. Gard. ser.* 2. *t.* 271.

A very pretty half hardy and half shrubby plant, native of Florida according to Nuttall, as quoted by Dr. Hooker, and of South Carolina according to Mr. Don.

Our specimens were furnished by Mr. Marshall, gardener to Mrs. Langley of Thames Ditton, in Oct. 1834. It is a neat, and very desirable plant, easily propagated by cuttings.

---

* Don Diego Gardoqui, Minister of Finance under Charles IV. of Spain, received the compliment of having this beautiful genus named after him, because of his having favoured the South American Expedition of Ruiz and Pavon.

1748

1748.



Miss Drake. del. Pub by J. Ridgway 169 Piccadi March. 1. 1835. S. Watts.

* VERBÉNA sulphúrea.

Sulphur-coloured Vervain.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. Verbenaceæ. *Juss.* (*Introduction to the Natural System of Botany, p.* 238.)

VERBENA.—Supra, vol. 4. *fol.* 294.

V. *sulphurea;* diffusa, pilosa, foliis 3-5-partitis: laciniis simplicibus lobatisque linearibus obtusis inferioribus ab imâ basi provenientibus, floribus subcapitatis, calycibus hispide pilosis, corollæ laciniis patentibus emarginatis.

V. sulphurea. *Don in Sweet's Flower Garden, ser.* 2. *t.* 221.

A pretty hardy perennial, with hairy stems spreading flat upon the ground, and heads of sulphur-coloured flowers.

Although not so conspicuous for its beauty as Verbena chamædrifolia (*V. Melindres* of this work, *t.* 1184.) it is nevertheless a neat pretty species, which deserves a place in all select collections.

It is a native of Chile, flowering in August, September, and November. Cuming found it near Valparaiso (No. 519); Bridges on the mountains near the same city. It is very nearly the same as *V. erinoides*, from which it differs in having yellow instead of purple flowers, and more particularly in the lower lobes of its leaves originating from the very base of the leaf, and not from near the point.

---

* See fol. 1184.

1740



Syd^s Edwards del.

Pub. by J. Ridgway 169 Piccadilly Ap. 1 1815

[illegible]

* DRACÆNA terminális.

The Sandwich Island Tee-Plant.

HEXANDRIA *MONOGYNIA.*

Nat. ord. Asphodeleæ, *Juss. (Introduction to the Natural System of Botany, p.* 271.)

DRACÆNA.—Supra, vol. 12. *fol.* 956.

D. *terminalis;* caule fruticoso vel arborescente, foliis petiolatis lanceolatis utrinque attenuatis, paniculæ ramis divaricatis simplicibus ramosisve floribus subsessilibus. *Römer & Schultes, Sp. Pl.* 7. 343.

D. terminalis. *Willd. Sp. Pl.* 2. 157.

Asparagus terminalis. *Linn. Sp. Pl.* 450.

One of the most graceful of arborescent stove-plants, where there is sufficient height for it to rear its slender stem to the elevation of ten or twelve feet. In appearance it resembles a Palm, and although, when its structure is carefully examined, it is found not to be one of that princely tribe, yet there can be no doubt that it serves to connect the chain of vegetation, by bringing the Asparagus in contact with the Cabbage Palm. Linnæus took it for a species of Asparagus.

In place of again describing a plant which has repeatedly been already in the hands of Botanists, we shall confine ourselves on this occasion to an extract from the entertaining work of Ellis on the Sandwich Islands.

"The *Ti* plant is common in all the South Sea Islands. It is a slow-growing plant, with a large, woody, fusiform

---

* See folio 1169.

1750

1750.



Miss Drake del.   Pub by J. Ridgway 169 Piccadilly Ap. 1. 1835.   S. Watts sc.

* BÉRBERIS dealbáta.

Whitened Barberry.

HEXANDRIA *MONOGYNIA.*

Nat. ord. **Berberideæ,** *Juss.* (*Introduction to the Natural System of Botany, p.* 30.)

BERBERIS.—Supra, vol. 12. *fol.* 1176.

---

§ *Folia simplicia, sempervirentia* (*coriacea*). *Flores racemosi.*

B. *dealbata;* subinermis, foliis simplicibus coriaceis suborbiculatis spinoso-dentatis subtus dealbatis, racemis oblongis densissimis pedunculatis.

Frutex *in horto 4-5-pedalis, verosimiliter orgyalis ad minimum, ramis purpureo-fuscis inermibus parcè foliatis.* Folia *coriacea, subrotundo-ovata, convexa, glaucescentia, spinoso-dentata, aliquandò subquadrata, imò cuneata 3-dentata, subtus albedine densâ corticata; petiolis brevibus basi articulatis.* Flores *in racemos oblongos multifloros densissimos nutantes congesti, lutei; sepalis apice sanguinolentis.* Petala *intus biglandulosa.*

A native of Mexico whence it was obtained by the Horticultural Society. Our drawing was made in the Chiswick Garden in April last.

This remarkable species is an evergreen shrub, which is probably hardy; but of this we are not certain, in consequence of its having been always hitherto protected in front of an east wall in winter. It differs in a striking manner from all other Barberries in several particulars. In the first place its leaves are very round and convex, and covered with a sort of bloom; then they are very white on the under surface, not however from the presence of any mealiness or white hairs, but on account of a discoloration of the cuticle; besides this there are scarcely any spines upon the branches; and finally its flowers are collected in remarkably dense nodding clusters, which are not much longer than the leaves. Its fruit is unknown.

---

* See folio 1176.

Like the other species of its genus it is most conveniently increased by layers, which will root in the course of a single season.

1751.



Miss Drake. del. Pub by J Ridgway 169 Piccadilly Ap.1 1835 J. Watts. sc.

1751

*LÆLIA ánceps.

Two-edged Lælia.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § EPIDENDREÆ, *Lindl.* (*Introduction to the Natural System of Botany*, p. 262.)

LÆLIA, *Lindl.* *Sepala* explanata, lanceolata, æqualia. *Petala* majora paulò difformia. *Labellum* (posticum), 3-partitum, lamellatum, circa columnam convolutum. *Columna* aptera carnosa, anticè canaliculata. *Anthera* 8-locularis. *Pollinia* 8, caudiculis 4 elasticis.——Herbæ *epiphytæ*, rhizomate *pseudobolbophoro*. Scapi *terminales, pauci v. multiflori*. Flores *speciosi, odorati.* Gen. & Sp. Orch. p. 115.

L. *anceps*; foliis binis aut solitariis lanceolatis, scapo ancipiti bifloro squamis carinatis vestito, ovario viscoso, labelli disco lineari elevato apice trilobo, pseudobolbis ovatis distantibus tetraquetris.

Rhizoma *repens, squamosum.* Pseudobulbi *ovati, distantes, tetraquetri, 2 poll. longi, juventute squamis acuminatis vestiti.* Folia *solitaria, rarò bina, lanceolata, acuminata, coriacea, atroviridia.* Scapus *ex apice pseudobulbi ortum ducens, sesquipedalis, gracilis, anceps, squamis carinatis arctè vaginantibus vestitus, apice biflorus.* Ovarium *viscosum, bracteâ lato-lanceolatâ, acuminatissimâ, carinatâ, membranaceâ, fuscâ, involvente brevius, apice teres, cuniculo a basi labelli orto brevi instructum.* Sepala *violacea, membranacea, dorso subherbacea, lanceolata, bipollicaria, patentissima.* Petala *oblongo-lanceolata, ejusdem coloris texturæ, et longitudinis, sed lineâ mediâ dorsi tantum herbacea, et duplò latiora.* Labellum *aut anticum, aut posticum, cucullatum, trilobum, basi columnam involvens et pallidè violaceum, intus luteum sanguineo-venosum; trilobum, lobis lateralibus rotundatis, intermedio atropurpureo basi albo, oblongo, acuto, subundulato, plano, lineâ latâ elevatâ crassâ luteâ antrorsum trilobâ in medio; lobis lateralibus abruptis, intermedio productiore in laminam abruptam verticalem attenuato.* Columna *semiteres, marginata, clavata, cum labello continua.* Anthera 8-*locularis.* Pollinia 8, *cuneata, utrâque extremitate caudicularum quatuor inserta et inflexa.*

A very beautiful Orchideous plant, imported from Mexico by Messrs. Loddiges, in whose collection our drawing was made last December. We at first took it for *Lælia grandiflora*, but Mr. George Loddiges has satisfied us that it is

not that species, which it is, however, highly probable that he possesses in a plant from Xalapa, with pear-shaped pseudo-bulbs, each of which has two or three lanceolate purplish leaves. The pseudo-bulbs of this are remarkable for their ovate outline, and for their four-angled figure, produced by a sharp ridge being planted upon each face of a compressed body.

When we say that this plant is equal in beauty to any of the Cattleyas, that it has a far more graceful mode of growth, in consequence of the length of its slender scaly stems, from the point of which the flowers swing, and that it diffuses an agreeable fragrance, we shall have said that it is one of the most interesting of the tribe that has yet made its appearance in our stoves.

It will probably succeed, without difficulty, in any hot-house which is adapted for the cultivation of Maxillarias and plants of that description.

1752

1752.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Ap. 1 1835. J. Watts sc.

* MONACHÁNTHUS víridis.

Green-flowered Cowlwort.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ § Vandeæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 262.)

MONACHANTHUS.—Supra, vol. 20. *fol.* 1735.

M. *viridis;* racemo multifloro, labello oblongo cuspidato margine lævi, sepalis petalisque rigidis ovatis.

M. viridis, *Lindl. Gen. & Sp. Orch. p.* 157.

Planta *facie omnino* Cataseti tridentati *quo foliis et caule tantum convenit ut floribus ademptis eandem diceres.* Flores *carnosi, herbacei, labelli margine vitellino.* Sepala *ovata, acuta, rigida, petalis parallela.* Petala *conformia, sed latiora et maculis quibusdam purpureis obscuris notata.* Labellum *posticum ventricosum, carnosum,* 1½-*poll. longum, ore obliquo lævi cuspidato.* Columna *brevis, ovata, carnosa, muticu.*

This is the original species on which the genus *Monachanthus* was founded. In habit it is so like *Catasetum tridentatum* that we long doubted whether it ought to be generically separated. The absence of the long tendril-like processes, which are so conspicuous upon the column of *Catasetum*, was, however, so remarkable a circumstance, that we came to the conclusion that it must be an essentially distinct form; and the discovery of the curious *Monachanthus discolor*, figured at tab. 1735, has shewn that we judged correctly.

A native of Brazil, growing upon trees in the Corcovado, whence Dr. Hooker received the drawing and specimen which first made the genus known. Our figure was made from a specimen communicated last November from Wentworth, by permission of Lord Fitzwilliam. We are not aware at what time, or by whom it was imported, but it has probably been taken for a green form of the common *Catasetum tridentatum*, and consequently no record has been kept of it.

---

* See fol. 1735.

It requires precisely the same treatment as *Catasetums*, and may be cultivated with facility by any one who has a damp stove.

1753

1753.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Apr. 1. 1835. S Watts sc.

* ARBUTUS procêra.

Tall Arbutus, or Strawberry Tree.

DECANDRIA *MONOGYNIA.*

Nat. ord. Ericaceæ, *Juss.* (*Introduction to the Natural System of Botany*, *p.* 182.)

ARBUTUS.—Supra, *vol.* 2. *fol.* 113.

A. *procera;* foliis oblongis serratis serrulatis integrisque glabris, petiolis calvis, racemis terminalibus paniculatis secundis.

A. procera, *Douglas Herb.*

Rami *juvenes glaucescentes, adulti rufo-castanei, glabri, salvis vegetissimis qui hispidi sunt; epidermide firmâ non deglubente.* Folia *in ramis surculosis oblonga, utrinque angustata, duplicato-serrata, petiolis hispidis; ramulorum oblonga, nunc cordata, integra vel irregulariter serrulata; petiolis ferè unciam longis.* Racemi *tomentosi, paniculati, terminales, secundi;* bracteis *ovatis, concavis, glabris.* Corollæ *albæ, leviter herbaceæ, ovatæ, medio paululùm constrictæ, basi obtusæ et obscurè decem-gibbosæ;* limbo *revoluto.* Fructus *juniores subrotundi, leviter obovati, tuberculati more Unedonis.*

A small tree found by the late unfortunate Douglas in the mountainous woody parts of the North West Coast of North America, and introduced by the Horticultural Society in 1825.

In the Gardens it forms an evergreen bush, with a fine broad glossy foliage, and a very vigorous appearance; but it requires to be carefully protected from extreme cold in winter, and succeeds best against a west wall. Its flowers are of a delicate greenish white, and are arranged after the manner of those of Arbutus Andrachne, to which species it is most nearly allied; differing however altogether in the form and serratures of its leaves, and in the form and size of its flowers.

---

* The Arbutus of Virgil and Horace was undoubtedly the modern Strawberry Tree. De Théis says the word has been formed from the Celtic *ar* rough, and *boise* a bush, in consequence of the astringency of its *fruit.* He considers that the term *horrida*, applied by Virgil to his Arbutus, has equally, in this case, the signification of austere.

The unripe fruits appear from our dried specimens to be like those of the common Arbutus, only obovate in a slight degree; the ripe fruit is unknown. The strong root-shoots are covered with scattered bristles, as also are the leaf-stalks, and the leaves themselves on such shoots are very strongly serrated.

Our drawing was made from a specimen obligingly supplied in May last, by Mr. Osborn, the present proprietor of the celebrated Fulham Nursery, lately occupied by Messrs. Whitley and Co.

1754

1754



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Ap. 1 1835 S. Watts sc.

1754

* BRÁSSIA Lanceána.

Mr. Lance's Brassia.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

BRASSIA.—Supra, vol. 10. *fol.* 832.

B. *Lanceana;* sepalis ovato-lanceolatis acuminatis, petalis minoribus, labello oblongo acuminato undulato sepalis lateralibus duplò breviore.

Color *totius herbæ amœnè et diffuso viridis.* Pseudobulbi *densi, oblongi, macri, compressissimi, ancipites, longitudinaliter sulcati, haud rarò arcuati.* Folia *bina aut solitaria, oblongo-lanceolata, macra, valdè striata.* Racemi *radicales, foliis longiores, floribus magnis luteis secundis odorem Primulæ suavissimum spirantibus onusti.* Bracteæ *ovatæ, concavæ, breves, membranaceæ.* Sepala *patentissima, lineari-lanceolata, purpureo paululum maculata, basi virescentia; supremum* 1½, *lateralia* 2 *poll. longa.* Petala *ejusdem formæ et coloris sed sepalo supremo duplo minora.* Labellum *luteum, immaculatum, oblongum, undulatum, ferè repandum, acuminatum, sepalis lateralibus duplò brevius; basi tuberculis* 2 *albis oblongis contiguis parallelis inanibus pubescentibus, dentibusque totidem membranaceis in fronte.*

A native of Surinam, in woods, where it was found growing upon trees by John Henry Lance, Esq.; and by him presented to the Horticultural Society in 1833. It is also wild in Brazil, where it was found by Dr. von Martius upon the trees on the banks of the river Jui, one of the tributaries of the Japurà in the Province of Rio Negro, flowering in February. It blossomed at nearly the same time in the Chiswick Garden, and with Messrs. Loddiges, in whose collection our drawing was made last August.

---

* Thus named by Mr. Brown in due commemoration of the late Mr. Brass, a skilful botanical traveller and draughtsman, who collected seeds, plants, and dried specimens on the Guinea Coast for Sir Joseph Banks, Dr. Fothergill and Dr. Pitcairn, and whose sketches being most liberally lent by Sir Joseph Banks to Dr. Afzelius in his visit to Sierra Leone, were maliciously damaged, and partly destroyed out of characteristic and wanton brutality by some piratical slave-mongers under the French flag, during the late war.—*Smith in Rees' Cyclopædia.*

It is of course, like other Surinam plants, a very tender species, and requires the hottest and dampest part of a stove; but there is no Orchideous plant which is more easy to cultivate, or more ready to multiply. It thrives in decayed leaf-mould, better than in any other compost, and may be readily known when out of flower, by the bright light green colour of its leaves, and by its peculiarly thin pseudobulbs, which often curve down upon themselves, as is represented in our figure.

If this is not equal to *Lælia* in the size of its flowers, and the brilliancy of its colours, it far surpasses that plant in its exquisite fragrance, which is like nothing so much as newly gathered cowslips and primroses.

1755

1755.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Ap. 1 1835. J Watts sc.

1755

[*] RHODOCHÍTON volúbile.

Twining Red-Cloak.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. SCROPHULARINEÆ, *Juss.* § ANTIRRHINEÆ (*Introduction to the Natural System of Botany, p.* 273.)

RHODOCHITON Zuccarini. Calyx membranaceus, coloratus, campanulatus, 5-fidus. *Corolla: tubus* anguloso-clavatus, internè pilis simplicibus reflexis, basi ubique, faucem versus 5-fariam vestitus; limbus 5-partitus, segmenta subæqualia erecta. *Stamina* didynama, rudimento quinti, erecta, apicibus simplicibus. *Stylus* sub stigmate rectus *Graham in Bot. Mag. t.* 3367. (*Capsula* vestita, pergamenea, bilocularis, polysperma, irregulariter dehiscens. *Semina* alata, radiatim striata; *ex Zuccarini.*)

Rhodochiton volubile; Zuccarini in litt. 1829, Otto et Dietr. in verhandl. ver. gart. Preuss. 10. 152. *t.* 1. Graham *l. c.*

Lophospermum atrosanguineum. *Zuccarini Plant. nov. et min. cogn. fasc.* 1. *in Abhandl. Math. Phys. Cl. Monac. vol.* 1. *p.* 306. *t.* 13.

Lophospermum Rhodochiton. *D. Don in Sweet's Brit. Fl. Gard. t.* 250.

A beautiful little climbing herbaceous plant, which was introduced late in 1833, by Mr. Low of Clapton, from the Berlin Garden. It is not very usual for us to be behindhand with our contemporaries in the publication of new plants, which are of much Horticultural interest; but on the present occasion we find ourselves forestalled in every direction. We console ourselves with the reflection that we are on that very account enabled to give a more complete history of the plant than would have been possible at an earlier period.

It was originally raised in the Botanic Garden at Munich from seeds collected in Mexico by Count Karwinski, and in the year 1829 was distributed to other gardens under the name of *Rhodochiton volubile*, an appellation which had been given it by Professor Zuccarini.

An account of it was soon after published in the Transactions of the Prussian Horticultural Society, by Messrs. Otto and Dietrich, with a tolerable figure, and the following account of the manner in which it had been treated in the Berlin Garden.

---

[*] From ῥοδος red, and χιτων a cloak, in allusion to the colour of its calyx.

"It is a climber, and may be applied to cover the walls in conservatories, as well as in the open ground during summer. It presents a really surprising aspect, when the outer walls of a greenhouse are covered with a screen formed by its numerous branches, tinged with red, and loaded with the most beautiful purplish-black flowers hanging down from the red calyxes. Out of every axil, to the most remote part of the twigs, a flower takes its origin, hanging down from a long stalk, which like the underside of the leaves and branches is coloured red. There can be no doubt, that it is, of all the known climbing plants with which we cover walls during summer, the most remarkable and beautiful, and ought to be particularly recommended, as it is easily cultivated, and flowers so very freely. Its propagation is by seeds and cuttings. The seeds are sown in March and April, that the young plants may become strong enough in time for planting in the open ground, where they remain during the whole summer, until frost sets in; if these plants are to be preserved, they should be potted off and put in a conservatory, where they prosper in a temperature, from 43° to 48° *Fahr.* They are also easily propagated by cuttings."

This mode of management appears to have been attended with great success, for when the plant was observed in the Berlin Garden in the autumn of 1833 by Dr. Henderson, it was in a much more thriving state than we have seen it in England.

Its next appearance in print was in the Transactions of the Mathematical and Physical Class (we believe) of the Munich Academy of Sciences, where it was published with an admirable description and figure by Professor Zuccarini; who, however, had altered his opinion of its belonging to a distinct genus, and called it *Lophospermum atrosanguineum;* this must have been in 1833, but our copy of the memoir to which we refer, bears no date.

Then Mr. Don, in August 1834, published it under the name of *Lophospermum Rhodochiton,* and he was followed in the succeeding December by Professor Graham, who adhered to Professor Zuccarini's first opinion, that the plant is distinct from Lophospermum, in which we entirely agree. In such groups as the Antirrhineous section of Scrophularineæ, less weighty reasons than the great coloured campanulate calyx, and salver-shaped corolla of *Rhodochiton,* as compared with the five-parted herbaceous calyx, and funnel-shaped corolla of *Lophospermum* are universally admitted as sufficient to justify the separation of such genera as *Antirrhinum, Linaria* and *Anarrhinum,* and we cannot conceive upon what principle the former are to be combined while the latter remain disunited.

In this country the *Rhodochiton* appears to require exactly the same treatment as *Lophospermum;* we saw it in great beauty at Mrs. Marryat's in a pot in the greenhouse last September, and it grew very well in the Garden of the Horticultural Society, trained to a pole in the open ground. Its greatest enemy seems to be bright sunlight.

1756

1756.



Miss Drake del.   Pub. by J. Ridgway 169 Piccadilly May 1.1835.   J Watts sc.

* DENDRÓBIUM Pierárdi.

Mr. Pierard's Dendrobium.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § MALAXIDEÆ, *Lindl.* (*Introduction to the Natural System of Botany*, p. 262.)
DENDROBIUM.—Supra, vol. 7. *fol.* 548.

D. *Pierardi;* caulibus pendulis glabris, foliis ovato-lanceolatis acutis, floribus geminatis racemum spurium formantibus, sepalis acuminatis membranaceis, petalis sepalo supremo majoribus acuminatis, labello cucullato dilatato subtruncato pubescente ciliato. *Gen. et Sp. Orch. p.* 79.
D. Pierardi. *Roxb. Fl. Ind.* 3. 482. *Hooker Exot. Fl. t.* 9.

Many years since an Orchideous Epiphyte was sent from Chittagong with some other species to the Botanical Garden, Calcutta, when Dr. Roxburgh named one of them *Dendrobium Pierardi*, in compliment to the gentleman who discovered it. In his Flora Indica Dr. Roxburgh adds that it is also a native of various parts of the Delta of the Ganges, where it is generally found on Mangoe Trees.

No one has however been able to discover any plant which answers exactly to Dr. Roxburgh's description of his *D. Pierardi;* and what is now cultivated under that name in the Calcutta Garden appears from Dr. Wallich's specimens to be the plant defined by Dr. Brown in the seventh volume of the first series of this work under the name of *D. cucullatum.*

When the first part of the *Genera and Species of Orchideous Plants* was published, we had satisfied ourselves that in all probability Dr. Roxburgh had confounded two or three different species under the common name of *D. Pierardi;* and accordingly we took the species represented by Dr. Hooker in his Exotic Flora, as the authority for the name, and we considered Dr. Brown's *Dendrobium cucullatum*, to be a mere form of it.

We have since had an opportunity of comparing *D. Pierardi* and *cucullatum* side by side, in the utmost perfection in the stove of the Messrs. Loddiges, where these lovely species flowered in great splendour in January last, forming festoons two or three feet long, quite covered with the most delicate pink and yellow blossoms. At first sight they are so entirely the same that one does not immediately perceive in what their differences consist; but upon a more attentive inspection it is found that *D. cucullatum* has larger flowers, with a broad roundish-ovate lip, the base of which is rolled up into a sort of short stalk, while in *D. Pierardi* the lip is rounded and very blunt, and its base is rolled into a much longer stalk, which quite conceals the column; it is moreover remarkably incurved. These differences are expressed at the bottom of the accompanying plate, where fig. 1. represents *D. Pierardi*, and fig. 2. *D. cucullatum*. Whether or not the latter is really a distinct species may possibly be doubted; but it is certainly so remarkable a form, as to have amply justified Dr. Brown in separating it from *Pierardi*.

---

* See folio 1239.

Allied to these plants are some new species, which we avail ourselves of the present opportunity of making known to Botanists.

1. D. *ochreatum* (Lindl. in Wall. Cat. no. 7410); caulibus brevibus e squamis membranaceis ventricosis erumpentibus, foliis ovato-lanceolatis acutis, vaginis ventricosis, floribus geminatis sepalis petalisque majoribus lanceolatis acuminatis, labello rhomboideo-ovato unguiculato cucullato intùs tomentoso.

Hab. in Chittagong, *Wallich* (*hab. s. sp. e Museo Anglo-Indico*).

Species vaginis laxis, alioquin facie D. Pierardi, primo intuitu distinguenda. Flores expansi ferè 3 uncias lati, pallidi. Labellum purpureo-maculatum; ungue intùs supra medium calloso.

A beautiful plant.

2. D. *Cunninghamii*; caulibus gracilibus pendulis ramosis vaginis foliorum corneis transversim corrugatis squamatis, foliis ovato-linearibus obtusiusculis, pedunculis oppositifoliis bifloris foliis multò brevioribus, sepalis ovatis acutis, petalis oblongis acutis latioribus, labelli trilobi lobo intermedio subrotundo undulato basi 5-lamellato lateralibus nanis acutis.

Hab. supra truncos Callistemonis elliptici (*A. C.*) in insula septentrionali *Novæ Zelandiæ*, juxta mare, incolis *Raumangha*; R. Cunningham (*hab. s. sp. comm. cel. A. Cunningham*).

This species is nearly allied to *D. biflorum*, of which I had examined no specimens at the time the *Gen. et Sp. Orch.* was published. Having since received that plant in a good state from Mr. Mathews, who found it in Otaheite hanging from the branches of trees, I am enabled to offer a correct definition of that very rare plant.

D. *biflorum* Swartz; caule pendulo gracili tereti, foliis lineari-lanceolatis acuminatis planis, pedunculis bifloris lateralibus e paleis corneis erumpentibus, sepalis petalisque acuminatissimis, labello rhomboideo medio bilamellato trilobo lobis lateralibus acutis nanis intermedio deltoideo acuminato margine fimbriato.

3. D. *Griffithianum*; caulibus erectis elongatis clavatis sursum tetragonis apice diphyllis, foliis lanceolato-oblongis obtusis, racemis laxis flexuosis multifloris pendulis e latere caulium natis, bracteis oblongis membranaceis pedicellis 4-plò brevioribus, petalis oblongis ciliolatis sepalis duplò latioribus, labello ovato pubescente denticulato subrepando, capsulis pyriformibus angulatis.

Hab. in Regno *Burmano* supra arbores, *W. Griffith*. (*hab. s. sp. comm. cel. Griffith*.)

A beautiful species with much the appearance of *D. aggregatum*.

4. D. *extinctorium*; pseudobulbis depressis orbiculatis aggregatis, foliis . . . ., pedunculis terminalibus unifloris, sepalis lateralibus posticis basi longè in extinctorii forma productis, labelli ungue longissimo tenui, limbo trilobo: lobis lateralibus erectis acutis intermedio oblongo rotundato, capsula pyriformi angulata.

Hab. in Regno *Burmano* locis humidis supra truncos Careyæ arboreæ, *W. Griffith* (*hab. s. sp. comm. cel. Griffith.*)

This is next *D. pusillum* Blume, and has very much the appearance of a Bolbophyllum. Its leaves have not been seen.

1757

1757.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly May. 1. 1836. J. Watts sc.

* BEGÓNIA petalódes.

Petaled Begonia.

MONŒCIA *POLYANDRIA.*

Nat. ord. BEGONIACEÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 169.)

BEGONIA.—Supra, vol. 4. *fol.* 284.

B. *petalodes;* caulescens, foliis æquilateris orbiculatis 5-9-lobis incisis serratis cucullatis, floribus masculis disepalis dipetalis, fœmineis tetrasepalis tetrapetalis, fructus alis subæqualibus acuminatis, cymis 2-3 floris.

Stipulæ *ovatæ, serratæ.* Foliorum *lamina petiolo brevior.* Cymæ *longè pedunculatæ* 2-3-*floræ.* Flores masculi *sepalis* 2 *subrotundis roseis, petalis* 2 *conformibus sed minoribus albis.* Flores fœminei *sepalis* 4, *parvis, subrotundis roseis, petalis totidem conformibus albis quorum* 2 *minora sunt.* Alæ *fructus immaturi æquales, acuminatæ.*

A pretty little species sent us by Mr. Richard Harrison, in April 1833, and we presume a native of Brazil. It adds another to the beautiful genus Begonia, which cultivators scarcely appreciate in this country. The species are all so very easily cultivated, and have so neat an appearance that they are exceedingly well adapted for ornamenting a damp stove. One of the prettiest things in the gardens of Schönbrunn is a Begonia house, filled with this genus and Ferns growing upon decayed wood and old tan.

To Botanists this is peculiarly instructive, because it shews better than any species we have previously seen what the regular form of the genus may be considered. To us it is the more interesting, because its petals, distinct from the sepals, shew we were right in considering the tendency of *Begonia* to be towards the production of a corolla, and that it is consequently correctly associated in the *Nixus plantarum* with Polypetalous orders; and secondly, that two or four being the number of the floral envelopes, when distinctly formed, the relationship of *Begoniaceæ* to *Onagrariæ* is almost demonstrated.

---

* See fol. 1252.

1758

1758.



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly May. 1. 1835. J. Watts sc.

* ONCÍDIUM citrínum.

Lemon-coloured Oncidium.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ § Vandeæ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

ONCIDIUM.—Supra, vol. 13. *fol.* 1050.

O. *citrinum;* pseudobulbis oblongis compressis, foliis ensiformibus rigidis scapo simplici brevioribus, sepalis petalisque labelli longitudine lineari-oblongis undulatis, labello cordato utrinque introrsùm arcuato apice dilatato subreniformi, cristâ 8-tuberculatâ pubescente, alis minimis, stigmate orbiculari.

Planta O. altissimo (*fol.* 1651.) *valdè affinis et fortè mera varietas. Diversa tamen videtur scapo non ramoso, floribus parum maculatis, sepalis petalisque minùs acuminatis, cristâ tuberculatâ potius quam digitatâ, demum alis minimis, et stigmate orbiculari nec angusto compressoque.*

A native of Trinidad, whence it was introduced by Messrs. Loddiges, in whose collection our drawing was made last November. Unfortunately the plant soon after sickened and died; so that it is for the present lost to the country.

It approaches very nearly to *O. altissimum*, figured at fol. 1651 of this work; and is principally distinguished by the following characters. Its flowering stem is simple and not branched; its flowers are of a pale lemon colour, very distant from each other, and by no means so much spotted; the crest of the lip consists of about eight warts, which are slightly downy, and not of nine smooth finger-like processes; its stigma is nearly orbicular, and not long and narrow, and the wings of the column are exceedingly small; and finally, both the pseudo-bulb and the leaves have a singularly yellow tint, as we are informed by Mr. George Loddiges.

Like the rest of the genus it requires a hot and damp stove.

---

* See folio 1542.

1759



Miss Drake del.   Pub by J. Ridgway 169 Piccadilly May 1 [illegible]   [illegible]

*PHYSIÁNTHUS albens.

White Bladderbloom.

PENTANDRIA *DIGYNIA.*

Nat. ord. ASCLEPIADEÆ, *R. Br.* (*Introduction to the Natural System of Botany*, *p.* 210.)

PHYSIANTHUS Martius. *Corolla* campanulata, tubo inflato-ventricoso, limbo 5-fido connivente. *Columna fructificationis* inclusa. *Corona staminea* inclusa, 5-phylla, foliolis tubo stamineo insertis, deinde corollæ adnatis, sursum liberis cucullatis. *Antheræ* membranâ terminatæ. *Pollinis* massæ 10, ceraceæ, compresso-clavatæ, in cruribus retinaculi deflexis pendulæ. *Stigma* bi-apiculatum. *Semina* comosa. *Martius nov. g. et sp.* 1. *p.* 53.

Ph. *albens ;* herbacea, volubilis, foliis oppositis integerrimis acutis basi cordato-truncatis subtùs albo pruinosis, floribus subdichotomo-cymosis. *Martius l. c. t.* 32. *Graham in Edinb. New Phil. Journ. Oct.* 1832. *Bot. Mag.* 3201.

Corolla *alba, coriacea, campanulata ; limbo patulo, laciniis ovatis undulatis medio roseo penicillatis ; tubo basi ventricoso.* Corona staminea *imâ basi corollæ inclusa ; appendiculis* 5, *ovatis, carnosis, extrorsùm cucullatis, inflexis, staminibus oppositis.* Stamina *connata in conum pistillum circumdantem ; antherarum loculis sagittatis, staminei coloris.* Pollinia *pendula, glandulâ nigrâ.* Stigma *globosum, apice bicorne.* Squamæ *hypogynæ O.*

We received a specimen of this rare plant in August last from Mr. Blair, gardener to Miss Martineau, as a native of Mexico. Upon comparing it, however, with the figure of *Physianthus albens*, given by von Martius from specimens collected by him in the woods of Ypanema in the Province of St. Pauls, and with the plate in the Botanical Magazine of the same plant from Buenos Ayres, we are obliged to conclude either that there has been some mistake with regard to its introduction to the London gardens, or that it stretches over a very great extent of the American Continent.

---

* Literally bladder-flower, in allusion to the inflated corolla.

It is a climbing plant, which runs rapidly over the rafters of a hothouse, or greenhouse, and may be easily propagated by cuttings.

1760

1760.



ss Drake del. Pub. by J. Ridgway 169 Piccadilly May 1. 1835. J. Watts sc.

* BLÉTIA refléxa.

Reflexed Bletia.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § EPIDENDREÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

BLETIA.—Supra, vol. 17. *fol.* 1401.

B. *reflexa ;* sepalis lineari-lanceolatis lateralibus reflexis, petalis cuneato-lanceolatis supra columnam conniventibus, labelli trilobi lobis lateralibus rotundatis planis intermedio angusto undulato lamellis 5 altis parallelis indivisis ad pedem columnæ decurrentibus, foliis angustis ensiformibus plicatis.

Folia *angusta, ensiformia, plicata.* Scapus 1½-2-*pedalis, teres, distanter vaginatus,* 2-4-*florus.* Bracteæ *oblongo-lineares, membranaceæ, ovario duplò breviores.* Sepala *angustò-lanceolata, ex viridi rubescentia basi alba ; lateralia reflexa, supremum paulò latius magis erectum, apice tantùm reflexum.* Petala *ex viridi rubescentia, cuneato-lanceolata, erecta, acuta, collateralia columnam a tergo tegentia.* Labellum *cucullatum, parte inferiore album, cum basi columnæ articulatum, oblongum, trilobum ; lobis lateralibus erectis rotundatis, albis, margine roseis, intermedio elongato pariter rotundato, crispo, atropurpureo, patente, lamellis* 5 *altis integris subundulatis totum axim labelli percurrentibus.* Columna *purpurea, clavata, apice alata, dente unico incurvo post cardinem antheræ.* Pollinia 8, *quorum* 4 *cæteris minora.*

Drawn in the hothouse of the Messrs. Loddiges, in November 1834. It is a terrestrial species, and a native of Mexico. The greenish colour of its flowers, and the remarkably narrow reflexed sepals, distinguish it at once from the remainder of the genus.

The specimen figured was less brilliant in its colours, and altogether smaller than is to be expected hereafter; for wild specimens now before us, collected by Baron de Karwinsky, and forming part of the Royal Bavarian Museum, have a rich purple cast upon every part of the flowers, and the scapes each bear from two to four blossoms.

Requires the same treatment as other Bletias.

---

* See folio 1401.

1761

1761.



Miss Drake del.   Pub. by J. Ridgway 169 Piccadilly May 1. 1835.   S. Watts sc.

* CAPRIFÓLIUM hispídulum.

Bristly Honeysuckle.

PENTANDRIA *MONOGYNIA.*

Nat. ord. CAPRIFOLIACEÆ, *Juss.* (*Introduction to the Natural System of Botany*, *p.* 206.)

CAPRIFOLIUM.—Supra, vol. 15. *fol.* 1232.

C. *hispidulum ;* tota hispido-pilosa, umbellis pedunculatis, corollis glabris bilabiatis tubo limbo duplò longiore, staminibus exsertis, foliis petiolatis cordato-ovatis obtusis subtus glaucis summis sessilibus liberis, caule filiformi.

Lonicera hispidula. *Douglas.*

Caulis *in genere debilis, filiformis, volubilis, vel prostratus, pilis rectis distantibus ut ferè omnes aliæ partes, hispidus.* Folia *parva.* Flores *parvi, rosei, pedunculis foliorum ferè longitudine, glomerulis bibracteatis.*

A very rare hardy shrub, discovered by Mr. Douglas in the woods of North West America. It is quite different from all the other honeysuckles, and is nearly scentless.

It will not readily grow more than two or three feet high, and seems as if it preferred lying prostrate to twining round other plants. In common soil it can scarcely be kept alive, but in peat and loam it grows as readily as any other hardy American plant.

Our drawing was made in the garden of the Horticultural Society in July last.

---

* See fol. 1232.

1762.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly May 1 1833. J. Watts sc.

1762

* LEPÁNTHES tridentâta.

Three-toothed Lepanthes.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § MALAXIDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

LEPANTHES Swz. Sepala patula, basi, præsertim lateralia, connata. *Petala* 2, libera, nana. *Labellum* apice sub-bilobum, cum columna connatum. *Columna* teres, nana, aut elongata. *Pollinia* 2.——Herbæ *epiphytæ,* caulibus *filiformibus, monophyllis, vaginatis,* spicis *terminalibus,* Pleurothallis *omninò habitu.* Flores *parvi, lutescentes vel rubescentes.*

L. *tridentata;* folio ovato acuto apice tridentato, pedunculis subbifloris folio longioribus, sepalis ovatis lateralibus semiconnatis, petalis semisagittatis tomentosis, labelli ciliati lobis lateralibus rotundatis, intermedio brevissimo acutiusculo incurvo, columnâ nanâ.

L. tridentata. *Swartz Prodr. p.* 125. *fl. Ind. occ. p.* 1561.

Caules *cæspitosi, vaginis lævibus laxis cincti, staturâ a lineis* 2 *usque ad uncias totidem variantes.* Folia *ovato-oblonga,* 1½ *unciam longa, crassa, apice acutè tridentata, subtus pallidiora.* Pedunculi *terminales, bracteati, subbiflori, foliis longiores.* Flores *parvi, bilabiati, oculo nudo purpureo-lutei.* Sepala *ovata, acuta, æqualia;* supremum *purpureum luteo-marginatum;* lateralia *semiconnata lutescentia, parum purpurata.* Petala *atropurpurea, tomentosa, semisagittata: lobo postico rotundato, sepalis dimidio breviore.* Labellum *minus, lutescens, basi concavum, purpurascens, cum basi columnæ connatum eâque parallelum, imò longius; trilobum, lobis lateralibus ascendentibus ciliatis columnam amplectantibus, intermedio incurvo minimo, minùs manifestè ciliato.* Columna *teres, minima, bicornis, basi tantum labello adnata; stigmate depresso subcuneiformi.* Pollinia 2, *parallela, conica, caudiculis apice incrassato aurantiaco cohærentibus.*

This which is one of the most pigmy of Orchideæ, not much exceeding in size the moss amongst which it grows, will nevertheless amply repay the observer for a careful examination of it. In order to explain its singular structure we have given magnified figures of the more important parts of its fructification.

---

* So called from λεπος bark, or λεπις small, and ανϑος a flower; because the plants of this genus have very small flowers and grow upon the bark of trees.

The first thing that strikes us is the singular termination of its leaves, which do not end in a sharp point as it seems to the naked eye; but have three little delicate teeth (fig. 2.). The flowers are covered externally by three sepals, the uppermost of which is purple with a yellowish border, and the lateral ones yellow with a little stain of purple; these are frosted over with brilliant little tubercles, and glitter in the sunshine as if encrusted with emeralds. The petals are two deep purple downy parts (fig. 2.), standing upright in the centre of the blossom, and forming a sort of back for the column to rest against; the column itself (fig. 5.) is a minute two-horned body, which buries its head between the lobes of the labellum (fig. 3.); and the latter is a yellowish three-lobed downy organ, whose two lateral divisions stand up on each side of the column, in the form of tiny bucklers ready to defend it from aggression, while the centre lobe projects in the fashion of a little horn (fig. 3. and 4.).

This is the first species of the genus which has been seen alive in Europe. It is a native of the highest parts of the mountains of Jamaica, where it is found growing on the bark of trees, among mosses. It can only be preserved alive with great care by being kept under a bell glass among damp moss in a cool part of a stove. We received it from Messrs. Loddiges in January last.

1763

1763.



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly May 1. 1835. J. Watts. sc.

* ÓROBUS atropurpúreus.

Dark Purple Orobus.

◆

DIADELPHIA *DECANDRIA.*

Nat. ord. LEGUMINOSÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 87.)

OROBUS L. *Calyx* campanulatus, 5-fidus, lobis 2 superioribus brevioribus. *Corolla* papilionacea. *Stamina* diadelpha. *Stylus* gracilis, linearis, apice villosus. *Legumen* cylindraceum, oblongum, 1-loculare, bivalve, polyspermum. *Semina* hilo lineari.——Herbæ *erectæ.* Stipulæ *semisagittatæ.* Petioli *in setum brevem simplicem desinentes.* Folia *abruptè pinnata, paucijuga.* Racemi *axillares, pedunculati.* De Cand. prodr. 2. 376.

O. *atropurpureus;* caule sub-simplici striato, foliis 1-plurijugis, foliolis linearibus acuminatis glabris, stipulis semisagittatis subunidentatis, pedunculis foliis longioribus, racemis densis secundis multifloris, corollis elongatis.

O. atropurpureus. *Desf. fl. atl.* 2. 157. *t.* 196. *De Cand. prodr.* 2. 386.

O. siculus. *Rafin. caratt.* 72.

O. Rafinesquii. *Presl. del. Prag.* 41.

A native of wild places near Algiers, where it was first noticed by Desfontaines. It is also met with in Sicily, and in the loamy meadows of eastern Calabria, near Cotrone and Cassano. In the gardens it is a hardy perennial, flowering in May; our drawing was made in the Garden of the Horticultural Society.

It varies in the number of its leaflets from one to several pair; Desfontaines' figure and ours represent extreme states.

---

* Said to be so named from ορω to excite and βους a bullock; that is to say, exciting to cattle. The Orobus of the Greeks was some sort of vetch.

1764

1764



J. G. Hart del.

1764

*CLAVÍJA ornáta.

Elegant Clavija.

PENTANDRIA *MONOGYNIA.*

Nat. ord. ARDISIACEÆ § C. THEOPHRASTEA, *Bartl. ord. nat. p.* 165. THEOPHRASTEACEÆ, *D. Don in G. Don. gen. syst.* 4. *p.* 24.

CLAVIJA.—Ruiz et Pavon. Calyx 5-partitus. *Corolla* rotata, 5-loba. *Coronæ appendices* 5. carnosæ, adnatæ. *Filamenta* in tubum connata! *Antheræ* cohærentes. *Ovarium* 1-loculare, multi-ovulatum. *Bacca* crustacea, mono-polysperma.——Arbores (Amer. Æquin.) *sempervirentes, trunco simplicissimo, recto, more* Palmarum *apice frondoso.* Folia *alternatim conferta, undique patentia, oblonga, coriacea, glabra, reticulato-venosa, margine dentato-spinosa v. integerrima;* petioli *basi callosi.* Flores *racemosi, albi aut aurantiaci. D. Don. MSS.*

C. *ornata,* foliis elongato-lanceolatis mucronatis dentato-spinosis, petiolis unguicularibus, racemis glabris, baccis subdispermis. *D. Don.*

Clavija ornata. *D. Don in edinb. phil. journ. January* 1831. *in G. Don gen. syst.* 4. *p.* 25.

Theophrasta longifolia. *Jacq. coll.* 4. *p.* 136. *hort. schœnbr.* 1. *t.* 116. *Spreng. syst.* 1. *p.* 670.

Arbor (*in caldario culta*) *vix ultrà* 6-*pedalis, tamen in loco natali sæpè altitudinem* 20 *pedum attingens, trunco recto, simplicissimo, apice frondoso, cortice fusco rugoso-rimoso induto.* Folia *conferta, subverticillata, patentia, oblongo-spathulata v. lanceolata, mucronata, dentato-spinosa, lætè viridia, glaberrima, cartilaginea, suprà nitidula, pedalia v. sesquipedalia, spithamam lata, costâ validâ pallidiori subtùs elevatâ, ramis primariis transversis parallelis, venulisque ramosissimis reticulata, basi attenuata, integerrima.* Petioli *unguiculares, robusti, fusco-purpurei, suprà planiusculi.* Racemi *supra-axillares, simplices, multiflori, palmares v. spithamæi, plerumque penduli, axi tereti, glabro, viridi.* Flores *odore* Primulæ veris *fragrantissimi, crocei, undique sparsi, inferioribus sæpè abortivis.* Pedicelli *brevissimi, robusti, teretes, glabri, pallidè aurantiaci.* Calyx 5-*partitus, persistens: laciniis rotundatis, coriaceis, impressè punctatis, margine membranaceis, leviter erosis, æstivatione imbricatis.* Corolla *rotata, substantiâ crassâ coriaceâ;* tubo *brevissimo, subobconico, intùs sulcato;* limbo 5-*lobo, patulo;* lobis *subrotundis, integerrimis, æstivatione imbricatis, disco convexiusculis.* Coronæ appendices 5, *flavæ, carnosæ, deltoideæ, adnatæ, tuberculiformes, suprà bisulcatæ, lobis corollæ alternæ.* Stamina 5, *corollæ lobis opposita;* filamenta *in tubum membranaceum infernè ventricosum connata;* antheræ *cohærentes, extrorsæ, biloculares, appendiculâ brevissimâ rotundatâ membranaceâ coronatæ;* loculis *oblongis, distinctis, parallelis.* Pollen *pulvereum.* Ovarium *liberum, pyramidatum, uniloculare, aurantiacum;* ovulis *numerosis,*

---

* Named by the authors of the Flora Peruviana in compliment to Jose Clavijo Faxardo, a Spanish Naturalist, who translated the works of Buffon into his own language.

placenta centrali insertis. Stylus *vix ullus.* Stigma *peltatum, 5-angulum pruinosum.* Bacca *magnitudine* Cerasi minoris, *subglobosa, crustacea, lateribus parùm compressa, unilocularis, evalvis, plerumque disperma, rarò mono v. trisperma, basi calyce et apice stylo persistentibus instructa,* crustâ *intùs virenti nervis venisque plurimis peragratá.* Semina *subrotunda, substantiá pulposá copiosè tecta,* subinde drupacea! aquâ tepidâ madida pulpa exit voluminosa, *basi umbilico amplo orbiculato concaviusculo instructa:* testa *cartilaginea;* albumen *magnum, corneum, pelliculá membranaceá arctè adhærenti vestitum.* Embryo *erectus, axilis, lacteus:* cotyledones *ovales, subfoliaceæ;* radicula *cylindracea, obtusa, cotyledonibus subæqualis, umbilico obversa. D. Don. MSS.*

This very elegant tree is a native of Caraccas, whence it was introduced to the Imperial Gardens at Schönbrunn, where it flowered for the first time in November 1789, and a description and figure have been given of it by the elder Jacquin in his magnificent work on the plants of that garden. A number of young plants were raised at the late Mr. Colvill's nursery, in 1829, from seeds collected in Caraccas by Mr. Fanning, to whom I am indebted for specimens of the ripe fruit. The young plants were observed to vary a good deal in the breadth of their leaves, and as some of them produced flowers in the spring of 1834, they were found to agree precisely with the plant here figured, which blossomed about the same time in the stove of the Chelsea Botanic Garden, and which had been previously introduced from the continent under the name of *Theophrasta latifolia.*

The plant loves a high temperature, and should be grown in a soil composed of equal parts of loam and vegetable earth, and plentifully supplied with water while in a growing state.

The late Sir James Edward Smith has expressed, in Rees's Cyclopædia, a suspicion of this species being identical with the *Eresia* of Plumier, the *Theophrasta americana* of Linnæus; but the latter having a large polyspermous fruit, and the former having seldom more than two seeds, completely sets the question at rest as to their specific difference at least.

The *Theophrasteaceæ*, consisting of *Theophrasta*, *Clavija*, *Jacquinia*, and *Leonia*, constitute a small group intermediate between *Myrsineæ* and *Sapoteæ*, being distinguished from the former by their polyspermous fruit, foliaceous embryo, extrorse anthers, and by the presence of coronary appendages alternating with the stamina, and which are to be regarded as the rudiments of a second series of those organs. *Theophrasta* agrees with the present genus in having extrorse anthers, but differs in its campanulate corolla, annular inclosed crown, and free stamina.

The dried flowers and seeds of *Clavija* and *Jacquinia* when immersed in warm water emit an agreeable fragrance resembling that of the Primrose. *D. Don.*

For these valuable notes we are indebted to Mr. Don, who had an opportunity of examining the specimen from which our drawing was made. We are only acquainted with the plant from the accounts of Jacquin to which Mr. Don has referred.

1765

1765.



Drake del. Pub by J. Ridgway 169 Piccadilly June 1. 1835. J. Watts sc.

* EPIDÉNDRUM grâcile.

Graceful Epidendrum.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 262.)

EPIDENDRUM.—Supra, vol. 1. *fol.* 17.

E. *gracile ;* foliis in pseudobulbos ovatos corrugatos pluribus levato-ensiformibus, racemo simplici longissimo, sepalis oblongis petalisque cuneatis patentibus, labelli ferè liberi trilobi lobis lateralibus semiovatis intermedio oblongo crispo obtusissimo duplò minoribus disco bicostato.

Labellum *tripartitum, cum basi columnæ leviter connatum; laciniis lateralibus erectis, semiovatis, obtusis, subundulatis, flavis sanguineo venosis, intermediâ subrotundâ crispâ ejusdem coloris brevioribus; in medio crassum, album venis purpureis, bicostatum; costis utrinque infra medium obsolete unidentatis.* Ovarium *brevè cuniculatum.* Columna *semiteres, marginata apice utrinque auriculata.* Gynizus *deorsum trilobus;* rostillum *crassum, breve, rotundatum.* Anthera *complete 4-loculares; dissepimentis membranâ marginatis.* Pollinia 4, *compressa basi unidentata; caudiculis totidem pulvereis replicatis.*

A native of the Bahamas, whence it was sent to the Horticultural Society by John Campbell Lees, Esq. in 1833. From a plant which flowered last September the accompanying drawing has been taken.

The plant is very nearly allied to *E. odoratissimum*, but its leaves are three times as long, and the flowering stem shoots up to the length of three feet, without a branch or a leaf beyond the little scales that appear upon it here and there; the flowers moreover are larger, and the form of the segments of the lip materially different, while the callosities upon the disk of the latter distinguish it positively. In *E. odoratissimum* the disk is simply thickened and flattened and gradually thins away towards the base ; in *E. gracile* it is distinctly two-ribbed, and has two obsolete teeth a little below the middle.

---

* See folio 1415.

The want of beauty in this plant is amply compensated for by its delicious fragrance, early in the morning and at night. It seems as if it were more terrestrial than epiphytal, and grows freely in any light well drained soil. The patch sent home by Mr. Lees consisted of a mass of pseudobulbs nearly two feet in diameter.

1766



Miss Drake del.   Pub. by J. Ridgway 169 Piccadilly June 1. 1835.   S. Watts sc.

1766

* VERBÉNA multífida; *var*. contracta.

Dwarf Purple Vervain.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. Verbenaceæ, *Juss. (Introduction to the Natural System of Botany, p.* 238.)

VERBENA.—Supra, vol. 4. *fol.* 294.

V. *multifida ;* diffusa pilosa, foliis 3-5-partitis laciniatis, corymbis subcapitatis pedunculatis, calycibus hispidè pilosis, corollæ laciniis patentibus emarginatis.
V. multifida. *Flora Peruv. vol.* 1. *p.* 21. *tab.* 33. *f. c. Syst. p.* 338. (1798.)
V. erinoides. *Willd. enum. p.* 634. *no.* 12. (1809!)—*Hooker Bot. Misc.* 1. 168.
Erinus laciniatus. *Linn. Sp. Pl.* 879.
"Lychnidea Verbenæ tenuifoliæ folio, vulgò Sandia Laguen. *Feuill. Per. v.* 3. *t.* 25."

One of the commonest of all plants on the alps of Chile and Mendoza, growing according to Dr. Gillies to the elevation of 8000 feet above the level of the sea; and varying extremely in the colour of the flowers, in stature, and in the degree in which the leaves are cut. In some individuals the flowers are said to be scarlet, in others blue, in the case before us purplish, and we can scarcely doubt that the *V. sulphurea* of fol. 1748 is in reality another form.

It is a hardy perennial; and one of its more robust varieties is now a common ornament of the flower garden, over which it runs during the summer, forming patches several feet in diameter.

What we now figure is a dwarfer and more short-jointed kind, our drawing of which was made in the Garden of the Horticultural Society last July. It looks almost like a species of scentless Thyme, and grows into a very dense patch which has but little disposition to extend itself.

---

* See folio 1184.

As there is no reason why the original name of *V. multifida* should be made to give way to Willdenow's alteration to *V. erinaides*, we retain the oldest designation.

1767



Miss Drake. del.   Pub by J. Ridgway 169 Piccadilly June. 1. 1835.   J. Watts.

1767

* GESNÉRA allagophýlla.

Shifting-leaved Gesnera.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. GESNEREÆ, *Richard.* (*Introduction to the Natural System of Botany*, p. 227.)

GESNERA.—Supra, vol. 4. *fol.* 329.

G. *allagophylla;* caule erecto piloso sæpè trigono, foliis villosis subsessilibus ternis oppositis vel sparsis lineari-oblongis spatulatisve obtusis crenatis, floribus sessilibus in verticillis spicâve longâ terminali, corollæ tubo piloso cylindrico: limbo æquali patenti glabro, glandulis hypogynis duabus posticis. *De Martius Nov. Gen. et Sp. pl.* 3. 36.

A stove plant, our drawing of which was made last July from a specimen belonging to Mr. Young of Epsom. Unfortunately we had no opportunity of making any notes concerning its structure, and consequently we are uncertain whether it is rightly referred to the above named species. It agrees very well with von Martius' careful description, except that we did not observe the upper part of the stem to be at all triangular.

If we are right in the species to which we have referred it, this is a native of the auriferous plains of Brazil in the Province of the Mines, in various places, especially about Villa Rica, the town of S. Joâo d'El Rey, in the Serras da Pires and da Piedade, and elsewhere. It has also been met with by Sellow, beyond the Tropic, in the Province of St. Paul's. *(Martius.)*

A very pretty neat species, and an interesting addition to the showy and easily cultivable genus to which it belongs.

---

* See folio 1158.

1768

1768.



Miss Drake del. Pub by J Ridgway 169 Piccadilly June 1. 1835. J. Watts sc.

* CAMPÁNULA Gargánica.

The Harebell of St. Angelo.

PENTANDRIA *MONOGYNIA.*

Nat. ord. CAMPANULACEÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 185.)

CAMPANULA.—Supra, vol. 1. *fol.* 56.

C. *garganica;* foliis radicalibus reniformibus longissimè petiolatis, caulinis cordatis omnibus crenato-dentatis pubescentibus, floribus axillaribus subfasciculatis (subsolitariis) corollis 5-partitis. *Alph. Decand. Monogr. des Campan. p.* 299.

C. garganica. *Tenore in Floræ neap. add. cum app. sem. h. r. neap.* 1827. *Sylloge, p.* 95. *Sweet's Brit. Fl. Gard. t.* 252.

For the opportunity of figuring this rare and beautiful species of harebell we are indebted to the kindness of Mrs. Marryat of Wimbledon, in whose rich collection it flowered in great perfection in July last. It had been raised from seeds presented to Mrs. Pallisser by Professor Tenore.

The following are the observations of that Botanist concerning this little known species. It is a perennial, flowering in June, and grows wild on Mount Gargano, among the ruins of the ancient Monastery, and near fort St. Angelo. It agrees in the form of its leaves, and in the branching of its peduncles with *C. Portenschlagiana;* but differs in the lobes of the calyx being twice not thrice as short as the corolla, and in the deep divisions of the latter. From *C. diffusa*, which it resembles in many respects, it may be distinguished by its deeply 5-parted corolla, with narrow lanceolate, revolute segments, and not half 5 cleft with broad ovate spreading lobes; the flowers moreover are racemose and not corymbose. From *C. Elatine* it differs in its stem being altogether diffuse and by no means ascending, tolerably firm, and from 3 to 6 inches long, in its root-leaves being nearly round and kidney-shaped, with double wavy crenellings, and not ovate, or between ovate and heart-shaped, with taper pointed pectinate teeth, in its equal corolla, which is pale blue and not bluish purple.

---

* See fol. 1738.

In many respects this account is somewhat at variance with the plant we have figured, and accords better with that published by Mr. Don in Sweet's Flower Garden; but our wild specimens from the shady rocks of Mount Gargano agree exactly with the individual that flowered in Mrs. Marryat's garden, and we conclude that the species is liable to variations in the crenelling of its leaves and in the colour and depth of the lobes of the corolla.

It is presumed that like most of the plants of the South of Italy this will require protection in winter from the heavy rains of this climate, otherwise it will probably prove hardy in the clefts of rockwork, of which it is a charming ornament.

1769

1769.



S. Drake del.   Pub. by J. Ridgway 169 Piccadilly June 1 1835.   J. Watts sc.

1769

*PSORÁLEA macrostáchya.

Long-spiked Psoralea.

DIADELPHIA *DECANDRIA.*

Nat. ord. LEGUMINOSÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 87.)

PSORALEA.—Supra, vol. 12. *fol.* 968.

P. *macrostachya;* foliis pinnatim 3-foliolatis pubescentibus, foliolis ovatis mucronatis, petiolis glanduloso-scabris, pedunculis axillaribus foliis quadruplò longioribus, spicis cylindraceis rachi bracteis calycibusque hirsutissimis. *Decand. prodr. v.* 2. 220.

Herba *perennis, erecta, ramosa,* 3*-pedalis, undique pubescens.* Foliola *integerrima, utrinque præsertim supra glandulosa; lateralia petiolulata.* Stipulæ *membranaceæ, ovatæ, acuminatæ.* Pili *racheos, bractearum, calycumque nigri densissimi; pedunculorum ferè nulli.* Flores *decidui, rachin hirsutum rectum linguentes.* Calycis *labium superius 4-dentatum.* Corolla *purpurea calyce parum longior;* alis *obtusissimis.* Legumen *ovatum, mucronatum, pilosum, calycis longitudine.* Semen *oblongum, aterrimum, glabrum.*

This species is described by De Candolle, from specimens communicated by La Gasca, and purporting to come from Nutka. Our drawing was however made from plants obtained from California through Mr. Douglas; and we therefore suspect some error in the former statement.

It is a handsome hardy perennial, yielding seed in tolerably plenty, and readily increasing by division. Our drawing was made in the Garden of the Horticultural Society in July of last year.

It is remarkable for the singular hairiness of the rachis and green parts of the flower, beginning abruptly with the origin of the first flower, but which is scarcely at all visible on the flower-stalk itself. After the flowers have fallen off this produces a singular appearance, the naked rachis remaining at the end of each flower-stalk in the form of a long hairy tail.

---

* So named, from ψωραλεος *warty*, in allusion to the very general presence of little tubercular secretions upon the surface of different species.

Fig. 1. represents a side view of an entire flower; 2. is a front view of the same; 3. a calyx with the corolla pulled off to shew the stamens; 4. a pod, and 5. a ripe seed.

177

1770.



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly June 1. 1835. Watts. sc.

* PENTSTÉMON staticifólius.

Sea Lavender-leaved Pentstemon.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. SCROPHULARINEÆ, *Juss.* (*Introduction to the Natural System of Botany*, *p.* 228.)

PENTSTEMON.—Supra, *vol.* 13. *fol.* 1121.

P. *staticifolius;* caule ascendente pubescente, foliis radicalibus oblongo-lanceolatis in petiolum longum angustatis integerrimis glabris: caulinis sessilibus cordato-ovatis dentatis pubescentibus, cymis subsessilibus calycibusque tomentosis, corollis ventricosis pubescentibus labiorum laciniis ovatis obtusis subæqualibus.

Caulis *subbipedalis, apice tomentosus et cymis subsessilibus verticillastros Labiatarum referentibus onustus.* Folia *radicalia cum petiolo 7-poll. longa.* Corolla *violacea, speciosa, fere pollicem et dimidium longa.*

A new hardy species of this beautiful genus, sent to the Horticultural Society by the late Mr. Douglas from California. It is most nearly related to *P. diffusus*, figured at fol. 1132 of this work, from which it differs in its much larger and more lilac flowers, in the form of its leaves, and in those next the root being perfectly entire.

Our drawing was made in the Garden of the Society in June last. At present the species is extremely rare; only one plant having been originally raised. It grows and flowers freely in a peat border.

---

* See fol. 1245.

We avail ourselves of a vacant leaf or two in this number to present our Botanical friends with a revision of the whole of the genera of the natural order to which Pentstemon belongs. For this valuable contribution we are indebted to George Bentham, Esq. the learned author of the Genera and Species Labiatarum.

Tribus I. **Verbasceæ**, *Bartl.* Corolla tubo abbreviato vel subgloboso, limbo explanato 4-5-fido vel bilabiato non ringente. Stamina fertilia 2-5, sæpius declinata. Antheræ approximatæ vel cohærentes nunc uniloculares nunc subbiloculares loculis divaricatis confluentibus. Capsula bivalvis septicide dehiscens, valvulis sæpe bifidis, dissepimento e marginibus valvulorum inflexis duplicato solubili. Semina nuda. Genera Solanaceis rotatis affinia sed æstivatione corollæ abunde distincta.

* *Corolla subrotata 5-loba.*

1. **Verbascum**, *Linn.* Stamina fertilia 5.
2. **Celsia**, *Linn.* Stamina fertilia 4. Capsula globosa usque ad basin dehiscens.
3. **Alonsoa**, *Ruiz et Pav.* Stamina fertilia 4. Capsula ovato-oblonga vix ad medium dehiscens. Flores torsione pedicelli resupinati. (Hemimeris *Auct. quoad species Americana*).

** *Corolla bilabiata labiis concavis.*

4. **Jovellana**, *Cav.* Corollæ lobi concavi subæquales patentes.
5. **Calceolaria**, *Linn.* Corollæ lobus inferior incurvus calceolatus.

*** *Corolla tubo subgloboso limbo valde inæquali.*

6. **Scrophularia**, *Linn.*

Tribus II. **Hemimerideæ.** Corolla tubo abbreviato, fauce concava, limbo explanato lacinia infima (sæpe bifida) basi concava 1-2-saccata vel calcarata. Stamina fertilia 2 vel 4 didynama adscendentia. Antheræ approximatæ uniloculares vel biloculares loculis divaricatis. Capsula bivalvis, loculicide dehiscens valvulis integris bifidisve vel septicide dehiscens valvulis bifidis, rarius indehiscens, dissepimento e marginibus valvularum inflexis duplicato solubili vel rarius concreto.

* *Corolla ecalcarata breviter bisaccata. Capsula globosa.*

7. **Angelonia**, *Humb. et Kunth.* Capsula loculicide dehiscens valvulis integris.
8. **Phylacanthus**, *Nees et Mart.* Capsula indehiscens.
9. **Hemimeris**, *Linn.* Capsula loculicide dehiscens septicide bipartibilis vel dehiscens. (Diascia, *Link.*)

** *Corolla calcarata. Capsula compressa apice truncata.*

10. **Nemesia**, *Vent.*

Tribus III. **Antirrhineæ**, *Chavannes.* Corolla tubulosa limbo personato vel ringente bilabiato rarius subæquali. Stamina fertilia 4 didynama. Antheræ per paria approximatæ biloculares. Capsula dentibus seu valvulis pluribus operculisve circumscissis dehiscens vel irregulariter rumpens. Semina nuda vel testa arilliformi inclusa.

* *Corolla basi calcarata.*

11. **Anarrhinum**, *Desf.* Corolla bilabiata. Capsula operculis 2 valviformibus dehiscens.

12. **Linaria**, *Tourn.* Corolla personata. Capsula operculis circumscissis vel plurimis dentiformibus vel valviformibus dehiscens.

* *Corolla basi saccata gibba vel æqualis.*

13. **Antirrhinum**, *Linn.* Corolla personata. Capsula apice poris vel foraminibus 2-3 dehiscens.

14. **Maurandia**, *Jacq.* Corolla personata. Capsula apice dentibus 10 dehiscens (Usteria, *Cav.*).

15. **Galvesia**, *Juss.* Corolla bilabiata. Capsula sub apice irregulariter rumpens; tubus corollæ intus nudus (Agassizia, *Chavannes*).

16. **Lophospermum**, *Don.* Corolla bilabiata. Capsula sub apice irregulariter rumpens; tubus corollæ intus pilorum seriebus 2 percursus.

17. **Rhodochiton**, *Zucc.* Calyx amplus coloratus. Corollæ limbus subæqualiter 5-fidus. Capsula sub apice irregulariter rumpens.

Tribus IV. **Salpiglossideæ.** Corolla tubo elongato rarius abbreviato, limbo obliquo nunc irregulariter bilabiato nunc concavo vel subplano lobis 5 sæpe incisis. Stamina fertilia 2 vel 4 didynama declinata. Antheræ biloculares loculis apice demum sæpius confluentibus. Capsula 2-4-valvis valvulis septicide dehiscentibus bifidisve. Genera plura a Solanaceis capsularibus (præsertim a Petunia) nonnisi æstivatione corollæ imbricativa et staminibus 4 nec 5 differrunt.

* *Corolla basi supra gibba limbo valde irregulari.*

18. **Collinsia**, *Nutt.* Corolla 5-fida. Capsula globosa 4-valvis.

19. **Schizanthus**, *Ruiz et Pav.* Corolla multifida. Capsula oblonga.

** *Corolla infundibuliformis limbo parum inæquali.*

20. **Salpiglossis**, *Ruiz et Pav.* Corollæ tubus ampliatus subrectus, limbi explanati laciniæ emarginatæ vel bifidæ.

21. **Browallia**, *Linn.* Corollæ tubus tenuis rectus apice parum dilatatus, limbi lobis integerrimis emarginatisve.

22. **Franciscea**, *Pohl.* Corollæ tubus tenuis apice incurvus, limbi lobis integerrimis planis.

23. ? **Brunsfelsia**, *Linn.* Corollæ tubus elongatus rectus æqualis, limbi lobis integerrimis planis. Capsula baccata.

24. **Anthocercis**, *Labill.* Corollæ tubus ampliatus limbo subrecto.

Tribus V. **Digitaleæ.** Corolla tubulosa sæpius ventricosa limbo bilabiato. Stamina basi declinata apice sæpissime adscendentia, 4 fertilia didynama, quinto summo sterili vel nullo. Antheræ biloculares loculis demum sæpissime divaricatis confluentibus. Stigma simplex, vel in lobis brevibus styli 2 vix incrassata. Capsula dura bivalvis, valvulis septicide dehiscentibus sæpius bifidis vel bipartitis, dissepimento duplicato e marginibus valvularum inflexis demum bipartibili. Placentæ a dissepimento demum sæpius solutæ. Genera priora Bignoniaceis affinia sed abunde distincta. Tribus habitu magis quam characteribus a Gratioleis differt.

* *Stamen quintum præsens sterile.*

25. **Chelone**, *Linn.* Corollæ labium superius amplum concavum. Semina membranaceo-alata.

26. **Pentstemon**, *Lhér.* Corollæ laciniæ subplanæ. Semina nuda.

** *Staminis quinti vestigium nullum.*

27. **Russelia**, *Jacq.* Corollæ laciniæ vix inæquales.

28. **Digitalis**, *Linn.* Corolla declinata labio inferiore longiore.

29. **Isoplexis**, *Lindl.* Corolla incurva labio superiore longiore.

Tribus VI. **Gratioleæ.** Corolla tubulosa limbo bilabiato. Stamina fertilia 2 vel 4 adscendentia. Antheræ biloculares muticæ. Capsula 2-4-valvis septicide vel loculicide dehiscens valvulis integris bifidisve, placentæ dissepimento non bipartibili demum libero adnatæ. Semina nuda.

§ 1. *Antherarum loculi disjuncti.*

30. **Pterostigma**, *Benth.* Stamina superiora fertilia antherarum loculis disjunctis, inferiora sterilia vel dimidiata. Calyx sub 5-partitus lacinia suprema majore. (Adenosma villosum, A. macrophyllum et Stemodia capitata. *Benth. in Wall. Cat. n.* 3851, 3853 *et* 3926, *cum specia quarta nova ex China.*)

31. **Lindenbergia**, *Link. et Otto.* Stamina 4 fertilia loculis disjunctis. Calyx semi 5-fidus (Brachycoris, *Schrader*, Bovea, *Decaisne*).

32. **Stemodia**, *Linn.* Stamina 4 fertilia loculis disjunctis. Calyx 5-partitus (Modestia et Diamoste, *Cham. et Schlecht.* Cybbanthera, *Hamilt.*)

§ 2. *Antherarum loculi contigui. Capsulæ valvulæ* 4 *vel si* 2 *loculicide dehiscentes vel fissæ.*

* *Stamina fertilia* 4. *Calyx semi-5-fidus. Capsulæ valvularum margines septiferæ inflexæ.*

33. **Dodartia**, *Linn.* Calyx tubuloso-campanulatus dentibus rectis. Corollæ labium inferius basi papillosum bigibbosum. Herba rigida paucifoliata.

34. **Mazus**, *Lour.* Calyx campanulatus dentibus patentibus. Corollæ labium inferius basi papillosum bigibbosum. Herbæ humiles scapis subnudis Hornemannia, *Willd.*).

35. **Limnophila**, *Br.* Calyx tubuloso-campanulatus labio inferiore glabro laciniis planis. Herbæ uliginosæ.

** *Stamina fertilia* 4. *Calyx* 5-*partitus. Capsulæ valvularum margines septiferæ inflexæ.*

36. ? **Capraria**, *Linn.* Corolla vix bilabiata. Stigma obtusum vix incrassatum (C. biflora, *Linn.*).

37. **Morgania**, *Br.* Corolla bilabiata. Stigma bilamellatum.

*** *Stamina fertilia* 4. *Calyx tubulosus* 5-*dentatus. Capsulæ valvularum margines subplanæ.*

38. **Mimulus**, *Linn.* Capsula 2-valvis.

39. **Uvedalia**, *Br.* Capsula 4-valvis.

**** *Stamina fertilia* 4. *Calyx* 5-*partitus. Capsulæ valvularum margines planæ.*

40. **Herpestis**, *Gaertn.* Capsulæ valvulæ bipartitæ. Calyces laciniæ valde inæquales (Monniera *Mich.* Bramia *Lam.*).

41. **Sphærotheca**, *Cham. et Schlecht.* Capsulæ valvulæ bipartitæ. Calycis laciniæ subæquales.

42. **Matourea**, *Aubl.* Capsulæ valvulæ subintegræ loculicide dehiscentes. Calycis laciniæ parum inæquales (*Species americanæ* 4-5 *inclusa* Gratiola acuminata, *Elliott.*).

***** *Stamina fertilia* 2. *Calyx* 5-*partitus.*

43. **Gratiola**, *Linn.* Stamina superiora fertilia.

44. **Beyrichia**, *Cham. et Schlecht.* Stamina inferiora fertilia. Capsulæ valvulæ vix fissæ (Sp. 4, Brasilienses).

45. ? **Achetaria**, *Cham. et Schlecht.* Stamina inferiora fertilia. Capsulæ valvulæ bipartibiles.

* * * * * * *Stamina fertilia* 2. *Calyx* 3-5-*dentatus vel* 5-*fidus.*

46. **Dopatrium**, *Hamilt.* Corolla calyce multo longior fauce ampliata (Gratiola juncea, lobelioides et nudicaulis).

47. **Microcarpæa**, *Br.* Corolla calycem minutum vix excedens.

§ 3. *Antherarum loculi contigui. Capsulæ valvulæ* 2 *integræ marginibus planis, dissepimento membranaceo parallelo.*

* *Stamina fertilia* 2.

48. **Peplidium**, *Delil.* Stamina sterilia nulla. Capsula sæpius irregulariter rupta valvulis vix solutis.

49. **Bonnaya**, *Link. et Ott.* Stamina sterilia 2. Capsula constanter valvatim dehiscens.

* * *Stamina fertilia* 4, *inferiorum filamenta basi appendiculata vel gibba.*

50. **Vandellia**, *Linn.* Calyx 5-partitus vel regulariter 5-dentatus non plicatus. Corollæ tubus intus nudus (Tittmannia, *Reichb.* Torenia, *Cham. et Schlecht.*).

51. **Artanema**, *Don.* Calyx 5-partitus. Corollæ tubus intus 4-squamatus (Diceros, *Pers. non Lour.*).

52. **Torenia**, *Linn.* Calyx tubulosus plicatus oblique 5-dentatus (Nortenia, *Dup. Thou.*).

* * * *Stamina fertilia* 4, *filamentis nudis integris.*

53. ? **Heteranthia**, *Nees et Mart.* Antheræ filamentorum inferiorum magnæ stylum amplectentes (Vrolikia, *Spreng.*).

54. **Lindernia**, *Linn.* Antheræ conformes. Calyx 5-partitus.

55. **Limosella**, *Linn.* Antheræ conformes. Calyx 5-dentatus vel 5-fidus.

Tribus VII. **Buchnereæ.** Corolla tubo tenui, limbo subplano 4-5-fido, laciniis sæpe bifidis. Stamina fertilia 4 didynama adscendentia. Antheræ uniloculares vel loculis demum divaricatis confluentibus. Capsula 2-valvis valvulis integris bifidisve. Genera nonnulla Selagineis habitu affines.

* *Capsulæ valvulæ loculicido-dehiscentes elasticæ integræ.*

56. **Buchnera**, *Linn.* (Campuleia, *Dup. Thou. tubo apice incurvo.* Piripea, *Aubl. tubo apice recto.*)

* * *Capsulæ valvulæ bifidæ marginibus septiferis inflexis.*

57. **Nycterinia**, *Don.* Calyx oblongo-tubulosus 5-angulatus 5-dentatus. Corollæ laciniæ emarginatæ vel bifidæ (Erini Capenses).

58. **Erinus**, *Linn.* Calyx 5-partitus. Corollæ laciniæ emarginatæ (E. alpinus).

59. **Manulea**, *Linn.* Calyx 5-partitus. Corollæ laciniæ integerrimæ inæquales (Nemia, *Berg.*).

60. **Sutera**, *Roth.* Calyx 5-partitus. Corollæ laciniæ 5 brevissimæ integræ æquales vix patentes (S. glandulosa, *Roth.* et Capraria multifida, *Linn.* ?).

Tribus VIII. **Buddleieæ.** Corolla tubo recto, limbo plano patente æqualiter 4-fido. Stamina fertilia 4 subæqualia. Antheræ distantes biloculares. Capsula 2-valvis.

61. **Buddleia**, *Linn.*

Tribus IX. Teedieæ. Corolla tubulosa bilabiata vel subinfundibuliformis, limbo subæqualiter 5-fido. Stamina fertilia 4 tubo inclusa didynama vel subæqualia distantia. Antheræ biloculares. Fructus baccatus.

* *Bacca indehiscens.*

62. Leucocarpus, *Don.* Corolla bilabiata. Calyx 5-dentatus.

63. Teedia, *Rud.* Corollæ limbus subæqualis. Calyx 5-partitus.

* * *Capsula baccata 4-valvis.*

64. Hemiphragma, *Wall.*

Tribus X. Veroniceæ. Corolla rotata infundibuliformis vel rarius irregulariter bilabiata. Stamina fertilia nunc 4 (vel 6-7 ?) subæqualia exserta distantia nunc 2 (superiora) inferiorum vestigiis nullis. Stylus simplex stigmate tenui vel rarius incrassato subcapitato. Capsula 2-valvis loculicide dehiscens valvulis sæpissime septicide bifidis bipartitisve. Semina nuda vel testa laxa arilliformi inclusa.

* *Stamina* 4 (*vel* 6-7 ?). *Corolla* 5 (*vel* 6-7 ?) *fida.*

65. Ourisia, *Comm.* Corolla campanulata 5-fida. Stamina 4.

66. ? Disandra, *Linn.* Corolla rotata 6-8-fida. Stamina 5-7. (*An hujus ordinis?*)

67. Sibthorpia, *Linn.* Corolla rotata 5-fida. Stamina 4.

** *Stamina* 4. *Corolla* 4-*fida.*

68. Scoparia, *Linn.* Corolla rotata.

69. Geochorda, *Cham. et Schlecht.* Corolla infundibuliformis.

70. Picrorhiza, *Royle.* Corolla campanulata calyce brevior. (Veronica Lindleyana, *Wall.*)

* * * *Stamina* 2.

71. Veronica, *Linn* Corolla rotata vel infundibuliformis (Pæderota, *Linn.* Leptandra, *Nutt.* Diplophyllum, *Lehm.* Cochlidiospermum, *Reichb.* Omphalospora, *Bess.* Hebe, *Juss.* Aidelus, *Spreng.*).

72. Calorhabdos, *Benth.* Corolla tubulosa incurva ore oblique 2-3-fido (Veronica Brunoniana, *Wall.*).

73. Wulfenia, *Jacq.* Corolla tubulosa recta limbo subbilabiato. Calyx 4-5-partitus.

74. Gymnandra, *Pall.* Corolla tubulosa recta limbo subbilabiato. Calyx 2-partitus (Lagotis, *Gærtn.*)

Tribus XI. Gerardieæ. Corolla campanulata infundibuliformis vel tubulosa, limbo 5-fido laciniis rotundatis planis. Stamina fertilia 4 adscendentia. Antheræ approximatæ 2-loculares loculis discretis parallelis sæpe acuminatis. Stylus simplex, stigmate obtuso sæpius elongato. Capsula 2-valvis loculicide dehiscens valvulis integris bifidisve. Semina sæpissime testa membranacea laxa arilliformi inclusa.

* *Calyx 5-dentatus. Corolla infundibuliformis.*

75. Escobedia, *Ruiz et Pav.* Corolla tubus elongatus limbo amplo 5-fido.

76. Physocalyx, *Pohl.* Calyx inflatus. Corollæ limbus parvus æqualiter 5-fidus.

** *Calyx 5-dentatus vel 5-fidus. Corolla campanulata vel tubulosa.*

77. Virgularia, *Ruiz et Pav.* Calyx 5-dentatus. Corolla ample tubulosa.

78. Macranthera, *Nutt.* Calyx profunde 5-fidus. Corolla tubulosa vel incurva. Capsula acuminata.

79. Gerardia, *Linn.* Calyx 5-dentatus vel semi-5-fidus. Corolla campanulata. Capsula obtusa (Melasma, *Berg.?* Sopubia, *Hamilt.*).

80. ? **Seymeria**, *Pursh.* Calyx 5-partitus. Corolla rotato-campanulata. Capsula acuta (Afzelia, *Gmel.*)

81. **Glossostylis**, *Cham. et Schlecht.* Calyx 2-5-dentatus. Corolla oblique campanulata. Semina intra membranam lineari-cuneatam minima. (Species observatæ tres: 1. G. aspera *Cham. et Schlecht.* Brasiliensis, 2. Rhinanthus scaber *Thunb.?* Capensis, 3. Hymenospermum dentatum *Benth.* Avensis, et forsan quarta inter plantas Senegalenses Perrottetianas).

*** *Calyx compressus hinc fissus.*

82. **Centranthera**, *Br.* (Razumovia, *Spreng.?*)

Tribus XII. **Rhinantheæ.** Corollæ limbus bilabiatus labio superiore concavo galeato vel lineari, integro vel emarginato, inferiore patente trifido. Stamina fertilia 4 didynama (rarius 2) adscendentia. Antheræ biloculares loculis discretis parallelis sæpe acuminatis. Stylus simplex. Capsula 2-valvis loculicide dehiscens valvulis sæpissime integris. Seminum nunc testa laxa membranacea inclusa, nunc nuda.

* *Antherarum loculi tenues mutici.*

83. **Orthocarpus**, *Nutt.* Antherarum loculi inæquales. Calyx tubulosus vel campanulatus 4-fidus.

84. **Castilleja**, *Mut.* Antherarum loculi inæquales. Calyx compressus bifidus vel quadrifidus hinc fissus (Euchroma, *Nutt.*).

85. **Schwalbea**, *Linn.* Antherarum loculi æquales. Calyx compressus oblique 5-fidus.

** *Antherarum loculi crassi sæpius acuti vel aristati.*

86. **Lamourouxia**, *Humb. et Kunth.* Corolla tubulosa compressa ventricosa labio inferiore brevi. Calyx 4-5-fidus.

87. **Cymbaria**, *Linn.* Corolla tubulosa ventricosa. Calyx 10-partitus.

88. **Odontites**, *Stev.* Calyx campanulatus 4-fidus. Corolla profunde bilabiata labio superiore laciniisque inferioris subæquilongis integerrimis vel intermedia retusa. Antheræ oblongæ galeam superantes.

89. **Euphrasia**, *Linn.* Calyx campanulatus 4-fidus. Corollæ labium superius galeatum emarginatum inferius magnum patens laciniis emarginatis (Parentucellia, *Viv.?*)

90. **Siphonostegia**, *Benth.* Calyx infundibuliformis tubo elongato limbo 4-5-partito. Corollæ labium superius incurvum inferius brevius (Species unica Chinensis).

91. **Bartsia**, *Linn.* Calyx campanulatus vel tubulosus 4-fidus. Corollæ labium superius galeatum compressum inferioris laciniæ integerrimæ. Antheræ ovatæ galea breviores (Trixago, *Stev.*).

92. **Pedicularis**, *Linn.* Calyx 5-fidus vel inæqualiter 2-3-fidus. Corollæ galea elongata sæpius incurva.

93. **Rhinanthus**, *Linn.* Calyx compressus membranaceo-inflatus. Corollæ labium superius galeatum. Semina pauca ala membranacea cincta (Alectorolophus, *Stev.*).

94. **Elephas**, *Tourn.* Corollæ labium superius lineare proboscideum, Semina pauca striata (Rhinanthus, *Stev.*).

95. **Melampyrum**, *Linn.* Corollæ galea compressa lobis replicatis. Semina in loculo solitaria magna exalata.

Genus incertæ sedis.

96. **Tozzia**, *Linn.* Corolla Gratiolearum, stamina Gerardiearum, vel Rhinanthearum. Capsula abortu monosperma.

Genera mihi non satis nota. Conobea, *Aubl.* Gomara, Sanchesia, Xuarezia et Calytriplex, *Ruiz et Pav.* Leucophyllum, *Humb. et Kunth.* Dichroma, *Cav.* Curtia, *Cham. et Schlecht.* Lafuentia, *Lag.* Hemianthus, *Nutt.* Willichia, *Mut.* (Hydranthelium, *Humb.* et *Kunth.*). Aptosimum, *Burch.* Hydrotriche, *Zucc.* Diceros, Picria et Tripinna, *Lour.* Bœa, *Commers.* Palmstruckia, *Retz, fil.*

1771.



Miss Drake del.

Pub. by J. Ridgway 169 Piccadilly July. 1. 1835.

S. Watts sc.

PÆÓNIA Moutan; *lacera*.

Double-red curled Tree Pæony.

POLYANDRIA *MONOGYNIA.*

Nat. ord. Ranunculaceæ, *Juss. (Introduction to the Natural System of Botany, p.* 6.)

PÆONIA.—Supra, vol. 1. *fol.* 42.

Garden Variety.

Raised from the seed of Pæonia Moutan by Mr. William Hyland, Gardener to the Earl of Sandwich, at Hinchingbrook, near Huntingdon, and communicated to us by Michael Foster, Esq.

This very beautiful variety is strikingly different from the other Moutans in the bright rosy red of the petals, the innermost of which are very much cut and gashed, curled up, and distinctly bordered with a narrow edge of light carmine, which sets them off to great advantage, and gives the whole flower a peculiarly rich and finished appearance. Mr. Hyland informs us that the seed was sown in September, and did not germinate for eighteen months afterwards; and that he finds such to be the case with all the seeds he has sown of this kind of Pæony.

It flowered for the first time in April 1834, when the plant was only three years old. It may therefore be expected that the future blossoms will be still finer than what we have figured.

1772

1772.



1

2

3

Miss Drake del. Pub by J. Ridgway 169 Piccadilly July 1. 1835. J. Watts sc.

* ANGRǼCUM micranthum.

Small-flowered Angræcum.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ § Vandeæ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

ANGRÆCUM.—Supra, vol. 18. *fol.* 1522.

A. *micranthum;* caule brevi, foliis oblongis trinerviis apice obliquis, spicis secundis multifloris horizontalibus congestis foliis brevioribus, sepalis petalisque subæqualibus angustè ovatis apice patulis, labello conformi basi bilobo bidentato medio pubescente, calcare obtuso incurvo postico.

Caulis 2-*pollicaris.* Folia 1½-*p. longa,* 1½-*p. lata, plana, crassa.* Flores *minuti, albi.*

A very small and unattractive species, lately introduced by Messrs. Loddiges from Sierra Leone. It is quite distinct from any previously described, but retains the peculiar characters of the genus without deviation. It flowered in February.

It is curious that Angræcum should be so exclusively an African genus, that no certain species has yet been found beyond its continent or immediately dependent islands, although it evidently stretches across the whole of the tropical part of Africa itself. In addition to this from Sierra Leone, three or four other species are now in our gardens from the same place, but all of them equally inconspicuous with the present. Of these *A. clandestinum*, *teretifolium* and *distichum* have already flowered with Messrs. Loddiges. Fig. 1. represents a lip with the spur attached; 2. an entire flower; and 3, a pair of pollen masses with their gland.

---

* See folio 1522.

1773



Miss Drake del. Pub by J. Ridgway 169 Piccadilly July 1 1835. J. Watts sc.

1773

[*] RUSSÉLIA juncea.

Rushy Russelia.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. SCROPHULARINEÆ, *Juss.* Trib. 5. DIGITALEÆ, *Bentham.* (*Introduction to the Natural System of Botany, p.* 228.)

RUSSELIA, Jacq.—*Corolla* tubulosa recta limbo patente subbilabiata labio superiore emarginato vel bifido inferiore trifido laciniis omnibus subæqualibus planis. *Stamina* fertilia 4, basi declinata apice adscendentia, sterile nullum. *Antheræ* sub labio superiore approximatæ glabræ biloculares loculis demum divaricatis confluentibus. *Stylus* simplex stigmate integro parum incrassato. *Semina* nuda.——Frutices, ramis *sæpius pendulis glabris.* Folia *opposita dentata vel integra.* Flores *nunc in racemis cymisve axillaribus dense glomerati nunc laxe racemosi pedunculis elongatis subunifloris.* Corollæ *sæpius coccineæ.* Bentham MSS.

R. *juncea ;* ramis tetragonis erectis junceis, foliis minimis petiolatis ovatis subintegris, pedunculis filiformibus subbifloris.

R. juncea. *Zuccarini in litt.*

Caulis *plantæ 3-pedalis erectus, angulatus, viridis,* ramis *onustus gracillimis, tetragonis, pallidè viridibus, junioribus subaphyllis, nutantibus.* Folia *sæpe minima, semper parva, ramulis debilioribus majora, petiolata, ovata, acuta, subtiliter serrulato-ciliata, aliquando dente uno alterove munita.* Inflorescentia *primo adspectu paniculata, sed revera e ramulis racemosim dispositis, gracilibus, distantibus constans, quorum apice insident flos unus duove pedicellati et cernui.* Calyx *parvus sepalis 5 ovatis acutis imbricantibus.* Corolla *tubulosa, coccinea, unciam longa, sursum dilatata, limbo subæqualiter 5-fido ; laciniis obtusis duabus superioribus magis approximatis.* Stamina 4, *didynama.* Antheræ *glabræ oblongæ, apiculatæ, lobis parallelis.* Stigma *bilobum ; lobis oblongis planis.*

A very pretty Greenhouse or Stove plant found in Mexico (probably near Oaxaca), by Count Karwinski, and lately introduced to the gardens of this country from

---

[*] Named after Dr. Alexander Russel, an English Physician, who resided for some years at Aleppo, and published an account of his observations upon the Natural History of that place in the year 1756.

Berlin and Munich. We have specimens from Dr. von Martius under the name we have adopted, but we know not in what publication Professor Zuccarini has described the species.

It forms a tufted light green bush, about three feet high, almost destitute of leaves, but having a graceful appearance in consequence of the large number of its drooping slender twigs, which look like those of an Equisetum or a Casuarina, and which form a singular contrast with the handsome scarlet flowers that appear in abundance in the months of July and August.

Propagated very easily by cuttings.

We are indebted to Mr. Bentham for the amended character of the genus. Fig. 1. is a corolla cut open by dividing the lip through the middle; 2. is a stigma with its style; 3. a stamen.

177

1774



Miss Drake del.   Pub. by J. Ridgway 169 Piccadilly July 1. 1835.   J. Watts sc.

1774

* ERIÓGONUM compósitum.

Compound Eriogonum.

ENNEANDRIA *TRIGYNIA.*

Nat. ord. POLYGONEÆ, *Juss.* Trib. ERIOGONEÆ, *Benth. in Linn. Trans. ined.* (*Introduction to the Natural System of Botany*, p. 169.)

ERIOGONUM Michx.—*Involucrum* tubulosum campanulatum vel cyathiforme vix angulatum subæqualiter 6-dentatum, multiflorum. *Receptaculum* bracteolis intra pedicellos sæpe instructum. *Flores* hermaphroditi vel dioici. *Perianthia* exserta profunde 6-fida laciniis 3 exterioribus, 3 interioribus sæpe dissimilibus. *Stamina* 9 basi subcoalita. *Ovarium* triquetrum. *Styli* 3 filiformes. *Stigmata* terminalia tenuia vel parum incrassata capitata. *Achenium* triquetrum vel vix trialatum.——Herbæ suffruticesve *sæpissime lanuginosæ.* Folia *caulina alterna ad basin caulis approximata cæspitosa vel secus ramos fasciculata exstipulata,* petiolo *basi dilatato sæpius amplexicauli vel vaginante.* Pedunculi *vel* rami floriferi *terminales nunc simplices* involucro *solitario terminali sæpius 2-3-chotome vel umbellatim ramosi, involucris solitariis glomeratisve, intra ramificationes, secus ramos vel ad apices sessilibus pedicellatisve.* Bracteæ *vel* folia ramorum floralium *sub ramificationibus orta tot quot rami vel (ramis abortientibus) numerosiora nunc minuta squamæformia, nunc herbacea et foliis caulinis difformia, sæpissime sessilia, exstipulata vel stipulis scariosis aucta.* Bentham in Linn. Trans. ined.

§ 3. *Umbellata,* Benth. Folia omnia exstipulata. Genitalia pilosa, filamenta prope basin ovarii. Inflorescentia umbellata pedunculo (sæpe scapiformi) apice umbellam simplicem vel duplicem pluriradiatam nonnunquam in capitulum globosum contractam vel ad involucrum solitarium reductam gerente. *Bentham l. c.*

E. *compositum* (Dougl. MSS.) foliis ad basin caulis approximatis longe petiolatis ovatis basi rotundatis cordatisve supra demum glabratis subtus dense albolanatis, pedunculo longissimo nudo apice breviter biumbellato, involucris breviter pedicellatis campanulatis multifloris. *Bentham l. c.*

Habitu et foliis *E. latifolio* affine, inflorescentia et floribus ad hanc sectionem referendum. Petioli *basi vaginantes, extus villosissimi,* 2-4-*pollicares, limbo* 1-1½-*pollicari.* Pedunculus *scapiformis ultrapedalis, in exemplaribus cultis folio sæpe instructis.* Bracteæ *sub radiis umbellæ lineares, oblongæ, vel (in exempl. cultis) dilatato-ovatæ.* Radii *umbellæ* 1½-3-*pollicares, umbellularum vix semipollicares, villosuli.* Flores *majores quam in E.* latifolio. Genitalia *minus pilosa quam in præcedentibus.* Perianthia *post anthesin aucta, laciniæ interiores elongatæ obovatæ, exteriores breviores latiores, marginibus membranaceis crispis.* Bentham MSS.

---

* Ἔριον *wool,* and γόνυ *a joint;* in allusion to the woolliness of the stems.

We are indebted to Mr. Bentham for the foregoing extracts from an unpublished paper read before the Linnean Society in May of the present year, and comprising the characters of several new genera, and species of this curious section of Polygoneæ.

The subject of the present article was found by Douglas on the rocky gravelly banks of rivers in New Albion.

It is a hardy herbaceous plant, forming a patch of neat dark green leaves which throw up in abundance, in May and June, their woolly scapes terminated by masses of yellowish white flowers. It thrives equally in peat earth, or common soil, best in a damp situation, and may be increased by cuttings of its well ripened shoots, struck in peat and sand in an almost exhausted hotbed.

We have illustrated the structure of this rare genus in more detail than can usually be afforded in a work so cheap as this. 1. Is a magnified view of a flower as seen externally. 2. Is a section to show the connection of the stamens at their base, and how they are placed by threes in face of the inner segments of the calyx. 3. An ovary with its recurved styles. 4. A fruit enveloped in the permanent calyx. 5. A ripe nut. 6. A seed taken from out of the latter. 7. A section of a seed to shew the position of the embryo in the albumen.

1775

1775.



Miss Drake. del. Pub. by J. Ridgway 169 Piccadilly July. 1. 1835. J. Watts. sc.

1775

* CLÍANTHUS puníceus.

Crimson Glory-pea.

DIADELPHIA *DECANDRIA.*

Nat. ord. Leguminosæ § Papilionaceæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 86.)

CLIANTHUS, Soland. Mss. in mus. Britt.—*Calyx* latè campanulatus, subæqualis, 5-dentatus. *Vexillum* acuminatum, reflexum, alis parallelis longius; *carina* scaphiformis, vexillo alisque multò longior, omninò monopetala. *Stamina* manifestè perigyna, diadelpha, omnia fertilia. *Stylus* staminibus duplò longior, versus apicem hinc leviter barbatus, stigmate simplicissimo. *Legumen* pedicellatum, coriaceum, acuminatum, ventricosum, polyspermum, intus lanulosum, suturâ dorsali rectâ ventrali convexâ. *Semina* reniformia, funiculis longiusculis affixa. ——Suffrutices, Herbæve foliis *impari-pinnatis stipulatis,* floribus *speciosissimis racemosis.*

C. *puniceus;* suffruticosus diffusus glaber, foliolis alternis oblongis subemarginatis, racemis pendulis multifloris, calyce 5-dentato, legumine glabro.

C. puniceus, *Soland. MSS. in mus. Britt. All. Cunn. in Hort. Trans. vol.* 1. *n. s. p.* 521. *t.* 22.

Donia punicea. *Don's gen. syst. of gard.* 2. *p.* 468.

Suffrutex *diffusus, ramosus,* 2-3-*pedalis, omni pube orbatus nisi levissimâ in paginam inferiorem foliolorum juniorum et inflorescentiæ partes virides; ramis viridibus, parum lignosis.* Folia 8-*juga cum impare;* foliolis *oblongis obtusis subemarginatis alternis;* stipulis *viridibus ovatis, reflexis, foliolis multò minoribus.* Racemi *penduli, multiflori;* axi *flexuosâ;* bracteis *angustè ovatis reflexis pedicellis bracteolatis filiformibus multò brevioribus.* Calyx *semi-quinquefidus, dentibus acuminatis.* Vexillum *ovato-lanceolatum, acuminatum reflexum* 2-*p. longum, atro-sanguineum versus basin albo interrupte vittatum, posticè roseum.* Alæ *atro-sanguineæ, obtusæ* 1½-*p. longæ.* Carina *omninò monopetala, acuminata, fere* 3-*p. longa, sanguineo-aurantiaca, basi pallida.* Legumen *ferè* 3-*p. longum, atro-fuscum, venosum.* Semina *reniformia, fusca, atro-nebulosa.*

The only account that has yet been given of this most beautiful plant is in the last part of the Horticultural Transactions; where a figure is published from a specimen furnished by Mr. Levison Gower from his garden at Titsey, near Godstow. We are indebted to the same gentleman for the specimen from which our plate has been prepared.

It is a native of New Zealand, and, in the opinion of Mr. Allan Cunningham, probably to the southward of the Bay of Islands; and

---

* From *κλεῖος glory*, and *ανϑος a flower*, in reference to the noble aspect of the species of this beautiful genus.

perhaps, on the shores of the River Thames, or at Mercury Bay. Its seeds were sent by the Missionaries under the name of *Kowaingutukaka*, or *Parrot's-bill;* and were stated to be the produce of a large tree; but this was no doubt a mistake, as there is no reason to believe that the plant will grow more than three or four feet high.

We extract the following account of it from the Horticultural Transactions.

"When planted in a peat border in the open air, where it succeeds the best, it forms a half-herbaceous evergreen shrub, not very unlike an evergreen Vetch, or more correctly speaking, a scarlet Colutea, (*Sutherlandia frutescens*). Its leaves are smooth, pinnated, and of rather a succulent texture, consisting of about eight pairs and an odd one. The stem is entirely free from furrows or angles, and is about as thick as a goose's quill. The flowers grow in oval clusters, hanging down from the axils of the leaves upon the lateral branches; each flower is rather more than three inches from the tip of the standard to the tip of the keel; the petals are of a light bright rich crimson, without any mottling or marking; the standard, which is of an ovate-lanceolate figure, and much tapered to the point, is reflexed so as almost to lie back upon the calyx; the wings are very much shorter than the keel, the point of which is so much prolonged as to look like the beak of some bird, although it must be confessed not much like that of a parrot. The flowers are succeeded by brownish black pods, two inches and a half long, seated on a slender stipe, and convex on the upper instead of the lower edge: so that unless attention be paid to their manner of growth, it would seem as if the seeds grew to the lower instead of the upper edge. They are covered all over inside with a delicate cottony down, in which lie the small kidney-shaped seeds, of a dull yellowish-ochre colour, mottled with small dark brown blotches and speckles.

"From the trials that have been made of the proper mode of managing it, both by Mr. Gower and the Rev. John Coleman, by whom it was given to the former gentleman, it would appear that it succeeds best when treated as a hardy plant, and turned out into a peat border; for in such a situation it has now been two years in Mr. Gower's garden, and the plants continue to look very healthy, with a profusion of blossoms forming for next year. Kept in a greenhouse it was sickly, and did not flower in the hands of Mr. Gower's gardener; but Mr. Coleman succeeded in blossoming it in a large pot in the greenhouse, and in inducing it to ripen its pods, one of which is that here figured.

"Considering that the climate of New Zealand is in some places so much like that of England, that some species, such as *Edwardsia microphylla*, will bear the rigour of our winters, it is not improbable that this may also prove a hardy plant. If so its extraordinary beauty will render it one of the most valuable species that has been introduced of late years; and even if it should be no hardier than *Sutherlandia frutescens*, it will still form one of the most important and welcome of all the modern additions to our flower gardens."

We refer our readers to the observations upon the other species made by Mr. Allan Cunningham in the place whence the foregoing passage has been extracted, for further information concerning these remarkable plants; adding only, that in the first place, the supposed attachment of the seeds to the lower suture of the pod, as described by Mr. Don, is a mistake, their attachment being in *Clianthus* as in all other Leguminosæ, namely, to the upper suture; and in the second, that as we are not aware of any reason why Dr. Solander's well-known name, *Clianthus*, should have been changed to that of *Donia*, we have not felt authorized to sanction the innovation.

1776

1770.



[illegible] Ridgway 169 Piccadilly July 1. 1835

* COLLETIA horrida.

Bristling Colletia.

◆

PENTANDRIA *MONOGYNIA.*

Nat. ord. Rhamneæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 113.)

COLLETIA, Kunth.—*Calyx* campanulatus, membranaceus, coloratus. *Petala* O, v. minutissima, linearia. *Stamina* antheris ovatis bilocularibus v. reniformibus unilocularibus. *Discus* brevis, cupulæformis, fundo calycis adnatus. *Ovarium* liberum, triloculare. *Stylus* simplex, elongatus. *Fructus* basi calycis tubo persistente cinctus, tricoccus, dehiscens. *Semina* sessilia. *Ad. Brong. in Ann. sc. nat.* x. 365.

C. *horrida;* spinis rigidis simplicibus vel tri-multipartitis, pedunculis sub geminatis, calycibus ovato-oblongis, staminibus sessilibus.
C. horrida. *Ad. Brong. l. c. an. Vent.* ?
C. ferox. *Gill. & Hook. in Bot. Misc.* 1. 154. *t.* 44. *f.* B.

Frutex *ulicinus, atroviridis, rigidus, nudus aut leviter pubescens, aphyllus, vel in ramis vegetioribus parcè foliatus.* Rami spinescentes *nunc omnino simplices, nunc tripartiti, sæpe multipartiti, validi et rigidi.* Folia *dum adsunt, minuta, ovata, serrata, atroviridia, glabra.* Pedunculi *sæpiùs geminati, nunc aggregati, sub anthesi nutantes, postea rigidi, ascendentes.* Calyx *ovato-oblongus, laciniis erectis, demum supra discum annularem revolutum transverse separabilis et deciduus.* Stamina *omninò sessilia.* Ovarium *superum, triloculare, ovulo unico erecto cuique loculo.* Stylus 1. Stigmata 3, *subcapitata.*

A hardy evergreen shrub, with much the habit of a furze bush, like which it consists chiefly of strong spines of a deep green colour. The flowers, which are of a greenish white, stained with dull purple, are produced in some abundance at the ends of the shoots, and appear in May and June.

Leaves the plant has none when old; but its young branches are furnished with them in the form of opposite bright green sawed scales, at the base of each of which is a small stipule. They however speedily fall off, leaving the branches to act as leaves by the aid of their soft parenchyma with which they are clothed in the form of bark.

---

* Named by Commerson after one Mons. Collet, a French Botanist, who is said by Jussieu to have been a fierce opponent of the method of Tournefort.

A native of Chili and Mendoza, where it is probably common. It is by no means so handsome a species as the *C. spinosa* of the same country, whose branches are loaded with masses of white bloom which must have an exceedingly pretty effect.

It grows in common garden soil, and prefers a hot exposed dry situation, such as the foot of a south wall, without any kind of shade. We have not as yet heard of its having been propagated; but it is often raised from Chilian seeds, under the name of *Retanilla*.

1777

1777.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly July 1 1835. S. Watts sc.

*PHOLIDÓTA imbricáta.

Imbricated Pholidota.

◆

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ § Malaxideæ. (*Introduction to the Natural System of Botany, p.* 262.)

PHOLIDOTA.—Supra, vol. 14. *fol.* 1213.

P. *imbricata*; pseudobulbis oblongis corrugato-sulcatis obtusis, foliis solitariis oblongo-lanceolatis plicatis acutis, spicis pendulis foliorum longitudine, bracteis membranaceis concavis acutis, sepalis lateralibus ovatis carinatis, labello subgloboso cucullato: lobis lateralibus parvis erectis intermedio bilobo: laciniis rotundatis.

P. imbricata. *Lindl. in Hook. Exot. Fl. t.* 138. *Loddiges Bot. Cab. t.* 1934.

It appears that there are two different species confounded under the name of *Ph. imbricata;* namely, this one and that which is figured at fol. 1213 of this work.

So long a time has elapsed since we saw the latter in flower that we are scarcely able to point out any very satisfactory characters to distinguish them, although we have no difficulty in acceding to the opinion of cultivators that they are in fact entirely distinct. We therefore propose to call *Ph. pallida* the plant before figured by the name of *Ph. imbricata*, and to retain the latter designation for this, which was originally figured by Dr. Hooker in the Exotic Flora, and afterwards by Messrs. Loddiges in their Botanical Cabinet.

They are both natives of India, and possibly occupy similar regions; they principally differ in the following particulars. *P. pallida* has very round blunt bracts, white flowers, smaller leaves, and grows very weakly and unwillingly under ordinary circumstances. *P. imbricata* has pointed bracts, yellowish flowers with a dash of violet, very long strong leaves, and grows and flowers most freely.

---

* See fol. 1213.

Our specimens were communicated by Richard Harrison, Esq., in May, 1834.

1778

1778.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Aug. 1. 1835 J. Watts sc.

1778

* MYÁNTHUS barbâtus.

Bearded Flywort.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ §. VANDEÆ. (*Introduction to the Natural System of Botany, p.* 262.)

MYANTHUS.—Supra, vol. 1. *fol.* 1538.

M. *barbatus;* labello in pilis succulentis barbæformibus dissoluto basi supra unicorni.

Caules *vetusti fusiformi-cylindracei,* 4-5-*poll. longi, parum induviati, luteo-olivacei.* Folia *atroviridia, undulata, Cataseti, oblongo-lanceolata, versus basin angustata.* Racemus *radicalis, prostratus vel pendulus, flores* 9-10 *inversos, i. e. labello quoad axin superiore gerens.* Bracteæ *lineari-lanceolatæ membranaceæ, pedicellis purpurascentibus breviores.* Perianthium *bilabiatum; sepalis petalisque lineari-lanceolatis, herbaceis, fusco-sanguineo maculatis.* Sepalum *supremum, cum petalis in galeam agglutinatum;* lateralia *patentissima.* Labellum *cum columna subarticulatum, lineare, medio infractum et saccatum; margine in fila tenuia alba numerosissima dissoluto, cristam aviculæ cujusdam simulantia; cornu deorsum falcatum, candidum, basi superiore tridentatum gerens.* Columna *herbacea, fusco-sanguineo maculata, cornu apicis galeæ longitudine, cirrhis duabus rectis purpurascentibus apice pellucidis cornu labelli amplectentibus.* Glandula *maxima, cartilaginea, oblonga, stigmate cum vi elasticâ dissiliens, et convoluta.*

This curious new species is a native of Demerara, near the falls of Wapopekai on the Massarony river, some hundreds of miles from its mouth, where it was found by Mr. John Henchman growing in the clefts of trees in shady situations. It was first sent us in February last by Mr. Lowe, with whom it first flowered, but almost immediately after we received it from Chatsworth, and from the garden of Mr. Willmore, of Oldford, near Birmingham. The first specimens were pallid, as if they had suffered from want of light, but Mr. Willmore's were much darker. In the latter the sepals were of a deep green spotted with purple, the petals were a deep rich brownish red; and the lip itself, with its innumerable thread-like feelers, was tinged with red in all parts, except a strong white horn, which stands upon it near its base, curving downwards towards the little pit which occupies the centre of the lip.

---

* See folio 1721.

This plant has altogether the habit of Catasetum, and no doubt requires the same sort of management.

1779



Miss Drake del.

* DENDRÓBIUM cúpreum.

Copper-coloured Dendrobium.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ §. MALAXIDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

DENDROBIUM.—Supra, vol. 7. *fol.* 548.

D. *cupreum;* caulibus teretibus pendulis, foliis oblongis obtusis emarginatis, racemis lateralibus laxis multifloris, bracteis ovato-linearibus canaliculatis obtusis, sepalis ovatis obtusiusculis petalis oblongis explanatis obtusis minoribus, labello unguiculato cochleariformi obtusissimo extus villoso intus nudo.

D. cupreum. *Herb. MSS.*

A native of the East Indies, whence it was sent by Dr. Wallich to the Hon. and Rev. W. Herbert, about the year 1825. It flowered at Spofforth for the first time at Midsummer 1834, when the drawing was made by Mr. Herbert, which we have now the opportunity of publishing.

It has been suspected that there are two Dendrobia in the East Indies, which agree with each other in a sort of tessellated appearance of the flowers, and a remarkably inflated slipper-like hairy labellum; one of these, called *D. moschatum*, has pale orange coloured blossoms, and a lip with 5 crested veins near its base; the other, to which the name of *D. Calceolaria* has been applied by Dr. Hooker, has rose coloured flowers with a little yellow in the centre, and is said to be destitute of the crested veins of the lip. There is, moreover, a *D. clavatum* of Dr. Wallich's Indian Catalogue, the specimens that were distributed of which were too imperfect to be described in the Genera and Species of Orchideous plants; but whether this plant is one of the preceding, or something altogether distinct is unknown. Furthermore, the plant now represented belongs to the same little cluster, and appears as if it might form a fourth species, unless indeed it should prove to be this *D. clavatum.* We have had no opportunity of examining it, but Mr. Herbert describes it to us in such a manner as to shew that it must be different from *D. Calceolaria* at least. "Its flowers are of a pale copper colour, veined with a redder tinge, and have two brown red blotches inside the lip. The spike of ten large flowers all expanded simultaneously, and the progress was so rapid that only about a week or ten days elapsed between its shewing bud and bursting into bloom. Independently of the colour of the flowers this species differs from *D. Calceolaria* in not having such long shoots, and in having the leaves less attenuated and shorter. *D. Calceolaria* under the same treatment makes shoots above four feet long; this plant under three feet. It is curious that these Dendrobiums if they miss flowering put forth a young plant instead of a spike of flowers at the point of inflorescence, as *D. Calceolaria* is doing here now."

---

* See folio 1239.

1780



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly Aug. 1. 1835. J. Watts sc.

*LASTHÉNIA glabráta.

Smooth Lasthenia.

SYNGENESIA *POLYGAMIA SUPERFLUA.*

Nat. ord. Compositæ. (*Introduction to the Natural System of Botany, p.* 197.)

LASTHENIA, De Cand. MSS.—Involucrum monophyllum, multidentatum, ebracteatum. *Receptaculum* conicum, scrobiculatum. *Flosculi* radii fœminei ligulati; disci hermaphroditi infundibulares. *Antheræ* apice appendiculatæ, basi muticæ. *Stigmata* rhombeolanceolata ad angulos barbata. *Pappus* O. *Achenia* compressa, lævia, apice disco brevi semicylindraceo mucronata.

L. *glabrata,* glaberrima; foliis subdentatis, involucris turbinatis.

Herba *annua, insipida, diffusa, glabrata, imò lucida, foliis oppositis a basi sub-amplexicauli acuminatis, subsucculentis, nunc integerrimis, nunc dentatis.* Capitula *majora quam in L. californica, flosculis magis vitellinis.* Ligulæ *oblongæ bidentatæ.* Flosculi disci *valde glandulosi.* Achenia *atra lævigata.*

A new hardy annual, native of California, whence it was introduced by the Horticultural Society in 1834. It flowers in May and June, if sown very early in the spring or in the previous autumn, and forms a pretty gay mass of yellow in the beds of the flower garden. It seeds profusely.

The characters of the genus Lasthenia have not yet been published; the name is employed in the manuscripts, of Professor De Candolle for another species,† which is also in our gardens, but which is rather less ornamental than the present.

---

* An unexplained manuscript name of Professor De Candolle.

† L. *californica* (De Cand.) subpubescens; foliis integerrimis, capitulis basi umbonatis.

1781

1781.





2.



S. Drake del.   Pub. by J. Ridgway 169 Piccadilly Aug. 1. 1835.   J. Watts sc.

* ANGRÆCUM distichum.

Two-rowed Angræcum.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ § VANDEÆ. (*Introduction to the Natural System of Botany, p.* 262.)

ANGRÆCUM.—Supra, vol. 18. *fol.* 1522.

A. *distichum* ; caule imbricato, foliis distichis compressis recurvis obtusis supra canaliculàtis, floribus solitariis axillaribus pedunculis foliis subæqualibus, sepalis ovatis petalisque angustioribus secundis obtusis, labello postico oblongo concavo apice tridentato calcare tereti horizontali pedunculo breviore.

Caules 3-4 *pollices longi, atrovirides, foliis densissimè imbricati.* Flores *parvi, lactei, carnosi, inodori.* Labelli *dentes laterales rotundati intermedio recto acuto.*

A remarkably neat and pretty looking species, on account of its small closely packed deep bright green leaves and tufted stems, but having no beauty of a striking nature in its flowers.

It is a native of Sierra Leone, where it is found on the bark of trees. Imported by Messrs. Loddiges, in whose Nursery our drawing was made in October last. It appears to grow freely in the damp hot atmosphere which so many similar plants delight in.

---

* See folio 1522.

1782

1782.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Aug. 1. 1835. J. Watts sc.

1782

* DÝCKIA rariflóra.

Scattered-flowered Dyckia.

HEXANDRIA *TRIGYNIA.*

Nat. ord. BROMELIACEÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 256.)

DYCKIA.—Schultes fil. Calyx tripartitus; laciniis erectis concavis. *Corolla* urceolato-campanulata; petalis erectis rhomboideis carnosis basi nudis. *Stamina* basi monadelpha; *antheris* erectis. *Ovarium* superum, tripartibile, polyspermum; *stylis* tribus *stigmatibusque* totidem dilatatis complicatis fimbriatis; *ovula* disticha.——Folia *carnosa,* epidermide *crassâ corneâ paululum lepidotâ vestita, margine spinosa.*

D. *rariflora;* foliis lineari-lanceolatis recurvo-patentibus, spicâ rariflorâ, spathis scapi ciliolato-serrulatis, floralibus calycem acutiusculum subæquantibus. *Schultes fil. in Rom. et Sch. Syst. veg. v.* 7. *p.* 1195. *Graham in Jameson's Journal, July,* 1835, *p.* 202.

Introduced from the Berlin Garden by the Horticultural Society in 1833. This plant is a native of the Serra of Villa Rica, in Brazil, where it and two more species were discovered by the indefatigable travellers Spix and Martius. It flowers in June and propagates very slowly by offsets, after the manner of an Aloe, with which it agrees in many of its habits.

---

* Named in compliment to his Highness the Prince of Salm—Reifferscheid-Dyck, a great lover of Gardening, and one of the most liberal and intelligent of the noble patrons of science of the present day.

The dry stove seems to suit it, for there it produces its rich orange flowers in great perfection, and retains them in all their freshness and beauty for several weeks.

We do not understand upon what principle this genus is referred to the Linnæan *Hexandria Monogynia*, instead of *Trigynia*, for it unquestionably has 3 distinct styles; unless it is to be considered a new case of the necessity of understanding the natural affinities of plants in order to use the sexual system.

Few persons, when they look at the leaves of a plant, ever think of the curious internal mechanism by which all its vital actions are put and maintained in motion; and yet there is not in the whole range of the creation a more singular object than a leaf, nor one whose structure is a more admirable instance of design and forethought. The internal anatomy in this species is highly curious and very easily examined. It consists as usual of a quantity of cellular matter enclosed in a cuticle, but the arrangement of the parts, which is most uncommon, is probably connected with the habits of life of the species in its native wildernesses. The cuticle is hard and composed on the upper surface of three, and on the lower of five layers of extremely minute compact cubical cells. The leaf itself is plane above and convex below; corresponding with the convexity is a stratum of equal thickness of dodecahedral cells, which are green, and pierced towards their upper side by the parallel veins of the leaf; above this structure is a very thick plano-convex bed of hard prismatical cells, which are planted nearly perpendicularly below the cuticle; so that when the section of the leaf is viewed by the naked eye it appears, as is indicated in our figure, as if it consisted of a common thin channelled leaf, whose concavity is filled up by cartilaginous matter. It is also worthy of remark, that in the lower stratum the tissue below each vein is much more lax and cavernous than that which is interspersed between the veins. The stomates are small and imperfect, and occupy a double or triple line in each furrow, lying concealed among the scurfiness.

178

1783.



[illegible] del. Pub. by S. Curtis [illegible] Aug. 1 [illegible] [illegible]

1783

* EMPÉTRUM rûbrum.

Red Crowberry.

DIŒCIA *TRIANDRIA.*

Nat. ord. Empetreæ, *Nuttall. (Introduction to the Natural System of Botany, p.* 109.)

EMPETRUM, L. ♂ *Squamæ* imbricatæ, quarum tres interiores petaloideæ. *Stamina* 3. ♀ *Squamæ* maris. *Ovarium* 6-9-loculare; *ovulis* solitariis ascendentibus. *Stigma* sessile, radiis plicatis tot quot loculi. *Pericarpium* baccatum, seminibus osseis.

E. *rubrum;* ramulis foliisque margine lanulosis.

E. rubrum. *Willd. sp. pl.* 4. 713.

Suffrûtex *sempervirens, virgatus, procumbens, visu subcinereo.* Rami *purpurascentes, graciles, ramosi, lanâ laxâ intricatâ vestiti.* Folia *linearia, 3 lineas longa, petiolata, marginibus veris reflexis in medio dorsi attingentibus, spuriis lanâ ciliatis.* Flores *axillares, solitarii;* masculi *e squamis constantes imbricatis, circiter novem, quarum 3 interiores corollæ formam induunt et sub aspectu petalorum trium linearium cuneatorum, fusco-purpureorum, apice reflexorum se expandunt.* Stamina 3; antheris *oblongis 2-locularibus in filamenta debilia vacillantibus.* Rudimentum *ovarii.* Fœminei *maribus sunt similes, sed omninò anandri.* Ovarium *sphæricum,* 9-*loculare.* Stigma *sessile, fusco-purpureum, radiis novem plicatis stellam referentibus.*

A hardy evergreen shrub, lately introduced by Mr. Low of the Clapton Nursery, under the name of "the Cranberry of Staten Island."

It is a plant that is found all over the Southern point of South America, growing with *Pernettia mucronata* along the sandy coast, spreading over the stones, but especially thriving at the back of the low sandhills, by which the shore is often skirted. It is, however, by no means confined to this locality, for we have specimens of it collected at Conception by Macrae. According to Gaudichaud the red berries are pleasant to eat.

---

* From εν *in,* and πετρος *a stone,* in allusion to the rocky places it grows among.

It grows freely in peat among other plants of a similar kind, and appears perfectly hardy. We have hitherto seen only fertile individuals alive, but in all probability sterile ones exist somewhere in collections, and if so we may soon expect to see its fruit.

Although not striking in its appearance, it forms an interesting addition to the hardy shrubs of this country.

1784.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Aug. 1. 1835.

J. Watts. sc.

* EUTÓCA divăricata.

Straggling Eutoca.

PENTANDRIA *MONOGYNIA.*

Nat. ord. HYDROPHYLLEÆ, *R. Br.* (*Introduction to the Natural System of Botany, p.* 244.)

EUTOCA.—Supra, fol. 1180.

E. *divaricata;* caulibus dichotomo-divaricatis, foliis omnibus ovatis indivisis, placentis 12-20-ovulatis. *Bentham in Trans. Linn. soc. vol.* 17. *p.* 278.

Annua. Caulis *decumbens, fragilis, pilosus.* Folia *indivisa, pilosa, petiolata, oblonga, obtusa.* Pedunculi *axillares, foliis longiores.* Corolla *calyce parum longior, subcampanulata limbo plano; plicis decem tubi squamularum loco.* Stamina *exserta.* Ovarium *ovatum, hispidum; stylo filiformi hispido, apice furcato.* Capsula *calyce acuto inclusa, ovata, acuta, compressa, pilis rigidis hispida; unilocularis, bivalvis; valvulis intus nitidis medio placentam corrugàtam polyspermam gerentibus.* Semina *plurima abòrtiva; matura ovalia, compressa, subtriangularia, scrobiculata;* embryone *tereti in axi albuminis cornei.*

A Californian annual of no great beauty, flowering in May and June. Its seeds should be sown in August in a situation neither overshaded, nor too much exposed to the sun; the young plants should not be allowed to stand nearer than four or five inches, and they should be kept carefully cleared of weeds; if this is not attended to, the branches will become so weak and brittle that they will not bear the water. Taking these precautions the flowers will be obtained in the spring, before the summer heats set in, and the seeds will ripen just when the season has become too dry to sustain the vegetation of the species.

The light violet blossoms are pretty in bouquets, but are hardly of more than Botanical interest.

---

* See folio 1180.

17

1785.



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly Sept. 1. 1835. J. Watts sc.

*GESNÉRA fauciâlis.

Wide-mouthed Gesnera.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. GESNEREÆ, *Richard.* (*Introduction to the Natural System of Botany, p.* 227.)

GESNERA.—Suprà, fol. 4. *fol.* 329.

G. *faucialis;* herbacea, foliis sub-sessilibus cordatis oblongis acutis crenatis rugosis tomentosis, racemo terminali, bracteis ovatis acutis reflexis, corollis tomentosis labio superiore oblongo bilobo basi angustato inferiore minimo revoluto fauce latissimâ truncatâ.

Communicated in July 1834, by the Hon. and Rev. W. Herbert, who received it from Mr. Tate. It is no doubt a native of Brazil, and is nearly related to *G. Selloi*, from which, however, it differs in many respects, especially in the great depth of the throat of the corolla from back to front, and in the narrow base of the upper lip.

Mr. Herbert considers it by far the finest of the genus, but he can only distinguish it by the size, brilliancy, and paucity of its flowers, from *G. bulbosa,* though it eclipses it. It is equally hardy, and thrives in the greenhouse, if not damp in the winter.

---

* See folio 1158.

178

1780.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Sept. 1. 1835 J. Watts sc.

1786

* ERYTHRÓNIUM grandiflórum.

Large American Dog's-tooth Violet.

HEXANDRIA *MONOGYNIA.*

Nat. ord. LILIACEÆ, *Juss.* (*Introduction to the Natural System of Botany*, *p.* 279.)

ERYTHRONIUM, L.—*Perianthium* hexaphyllum, campanulatum, foliolis reflexis: petalis basi bituberculatis. *Stamina* hypogyna, tria longiora. *Stylus* trigonus. *Stigma* 3-lobum v. 3-sulcatum. *Capsula* polysperma, loculicido-trivalvis. *Semina* ovata.——Folia *radicalia*, *gemina;* flores *speciosi;* scapi *sæpius uniflori*, *nunc ramosi.*

E. *grandiflorum;* foliis oblongo-lanceolatis subcomplicatis obtusis, perianthii laciniis ovato-lanceolatis acuminatis fere ab imâ basi reflexis, stigmate tripartito.

E. grandiflorum. *Pursh. fl. am. sept.* 1. 231.

Species Erythroniis alteris americanis affinis sed valdè distincta. Ab americano *vulgari et* Nuttalliano *differt foliis concavis impunctatis et stigmate tripartito; ab* albido *formâ foliolorum perianthii diversissimâ et stigmate altiùs trifido; ab omnibus floribus duplò majoribus intensiùs luteis, foliolis ab imâ basi ferè reflexis.*

Of this extremely rare plant, a single bulb was received by the Horticultural Society from North West America eight or nine years ago; it has continued to grow slowly in a peat border, and at last put forth its beautiful flowers last May. It has not however produced any seed, and it will probably be many years before it can possibly be distributed.

Mr. Douglas, who discovered it, considered it the *Erythronium grandiflorum* of Pursh; and we adopt his opinion, although we confess ourselves unable to reconcile with it the statement by that writer, that his plant has linear-lanceolate leaves.

---

* The common Dog's-tooth violet was the *σατύριον ἐρυθρόνιον*, or *red Satyrion*, of Dioscorides, and hence the latter word has been adopted by the moderns.

Its very large flowers, with their segments bent back almost to their base, distinguish this at first sight from the other American species.

Beautiful as it is this cannot be compared with an unpublished species from the same country, which we possess from Mr. Douglas; and which is most remarkable for having *an irregularly branched scape*. This may be named and characterised as follows:—

Erythronium giganteum; foliis oblongis lanceolatisve, scapo irregulariter ramoso 5-floro foliis duplò longiore, foliolis perianthii acuminatis medio reflexis, stigmate 3-lobo.

1787

1787.



[illegible] Ridgway 169 Piccadilly Sept. 1 1835 [illegible] Watts

* ONCÍDIUM pulchellum.

Pretty Oncidium.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ, *Juss.* § Vandeæ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

ONCIDIUM.—Suprà, vol. 13. *fol.* 1050.

§. 2. Folia triquetra v. teretia.

O. *pulchellum;* foliis acutè triquetris carinatis subfalcatis integris, scapo confertissime multifloro subcernuo, sepalis cymbiformibus acuminatis lateralibus connatis, petalis ovatis undulatis acutis, labelli lobis lateralibus rotundatis intermedio subrotundo sessili retuso v. emarginato subæqualibus, cristâ trilobâ anticè 1-callosâ, alis columnæ acinaciformibus denticulatis.

O. pulchellum. *Hooker in Bot. Mag. t.* 2773. *Lindl. Gen. & Sp. Orch. p.* 206. *Lodd. Bot. Cab. t.* 1984.

A beautiful species found in several parts of the West Indies, and probably mistaken for *O. variegatum* till Dr. Hooker distinguished it. When in flower its panicle is so loaded with white blossoms tinged with yellow and pink as to be weighed down with their profusion.

It requires the treatment usually given to its tribe.

---

* See folio 1542.

178



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Sept. 1. 1835

S. Watts sc.

* AZARA dentáta.

Toothed Azara.

POLYANDRIA *MONOGYNIA.*

Nat. ord. BIXINEÆ, *Kunth.* (*Introduction to the Natural System of Botany, p.* 152.)

AZARA, Ruiz et Pav. Cal. 4-6-partitus. *Petala* O. *Stamina* plurima filamentis filiformibus liberis. *Antheræ* biloculares. *Ovarium* 1-loculare, placentis tribus polyspermis parietalibus; *stigma* indivisum. *Bacca* stylo apiculata, 1-locularis, polysperma.——Frutices *Chilenses.* Stipulæ *nullæ, v. aliquando solitariæ folium parvum mentientes.* Flores *fragrantes.*

A. *dentata;* ramis pubescentibus, foliis oblongis glabris nitidis crenato-serratis, stipulis rotundis, racemis corymbosis foliis brevioribus, staminibus calyce parum longioribus.

A. dentata. *Ruiz et Pavon. Fl. Peruv. syst. p.* 138.

A common shrub in the woods near Conception in Chile, also frequent near Valparaiso.

In this country it forms a very handsome evergreen bush, with remarkably glossy deep bright green leaves. It appears quite hardy if nailed to a wall and protected from wet in winter, but it will not bear the winters near London under other circumstances. No drought seems to affect it; for at the time we are writing, after nearly two months of the hottest and driest weather known in England, on the face of a south wall, the leaves of this plant are perfectly fresh and green.

It may be increased by cuttings, but layers are preferable.

---

* Named after Joseph Nicolas Azara, a Spanish Gentleman, of whom nothing further is known, except that he was a patron of science.

1789



[illegible] del.

Pub. by J. Ridgway 169 Piccadilly Sept. 1 1835.

S. Watts sc.

*ONCÍDIUM Lemoniânum.

Sir C. Lemon's Oncidium.

GYNANDRIA *MONANDRIA.*

Nat. Ord. ORCHIDEÆ, *Juss.* § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

ONCIDIUM.—Suprà, vol. 13. *fol.* 1050.

---

§ 2. Folia triquetra v. teretia.

O. *Lemonianum;* foliis compressis acuminatis suprà sulcatis, scapo stricto paucifloro, sepalis parvis spatulatis apiculatis omnibus liberis, petalis oblongis undulatis, labelli laciniis lateralibus linearibus abbreviatis intermediâ maximâ reniformi bilobâ: lobis dentatis ungue brevi margine denticulato, columnæ alis subquadratis truncatis oblique unidentatis.

"This curious little epiphyte was among a collection imported from the Havannah in March last, by Capt. Sutton of Flushing near Falmouth, and by him presented to Sir Charles Lemon, Bart., M. P., in whose garden at Carclew it flowered in May. It appears to us to be totally different from any species hitherto noticed, and is readily distinguished by its peculiar foliage."

"*Pseudobulbs* very small, scarcely discernible, roundish, one-leaved. *Leaves* from one to three inches long, and about a quarter of an inch broad, dagger-shaped, two-sided or edged, and sharp at the point; pale green, slightly striated, and covered with a profusion of minute, roughish, pale dots or warts. *Scape* about 9 inches high, round and slender, nearly erect, and in our specimen 5-flowered. Its colour is a pale brownish green, unless at the joints where it is a dark brown. *Sepals* yellowish, marked along the back with reddish spots. The upper one which rises erect is spathulate, and appears to have generally 2 spots on the inside behind the column. The two lower ones are small and acuminate. *Labellum* bright yellow, beautifully spotted. The lower division is crescent-shaped, two-lobed, slightly unguiculate and fringed at the outer edge. The upper divisions are panduriform and of the same colour as the lower, but without the spots. The centre is tuberculated and ciliated on each side. *Column* deep yellow, with comparatively large spreading wings."

---

* See folio 1542.

For the previous note we are indebted to Mr. Booth, under whose successful management this and several other curious new Orchideous plants have been raised at Carclew. The species is nearly allied to O. Cebolleta, from which, however, its characters abundantly distinguish it.

1790.



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly Sept. 1. 1835. J. Watts.

* KENNÉDYA Marrýattæ.

Mrs. Marryat's Kennedya.

DIADELPHIA *DECANDRIA.*

Nat. Ord. LEGUMINOSÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 86.)

KENNEDYA.—Suprà, vol. 11. *fol.* 944.

§ 1. *Foliis trifoliolatis, carinâ rectâ vexillo sublongiore,* De Cand.

K. *Marryattæ;* foliolis tribus oblongis obtusis undulatis petiolo brevioribus, junioribus caulibusque villosissimis, stipulis bracteisque cordatis apiculatis, pedunculis 4-floris.

A beautiful greenhouse climber, obtained from Swan River seeds sent home by Sir James Stirling. Our first specimens were communicated by Mrs. Marryat, we afterwards received others from the garden of Mr. Robert Mangles. It flowers abundantly from April to July, producing a striking appearance with its numerous scarlet blossoms.

As it is easily propagated by cuttings it will soon become a common plant.

Different as this is from *K. prostrata* it is not easy to fix upon any very decisive mark of distinction. It is much larger in all its parts, and very much more hairy; in fact its young leaves and stems are perfectly shaggy. The leaflets are larger, far more wavy, and shorter than their stalk. The flowers grow about four in a cluster, instead of two, or being altogether solitary. The fruit we have not seen.

---

* See folio 1421.

1791



Miss Drake. del.   Pub by J Ridgway 169 Piccadilly Sept. 1. 1835.   J Watts sc.

* ARCTOSTÁPHYLOS tomentósa.

Downy Bearberry.

DECANDRIA *MONOGYNIA.*

Nat. ord. Ericeæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 182.)

ARCTOSTAPHYLOS, Adanson. Omnia Arbuti, sed *Drupa* lævis, composita, putamine pluriloculari, loculis monospermis.——Frutices.

A. *tomentosa;* ramis tomentosis sæpius setoso-hispidis, foliis ovatis acutis coriaceis junioribus tomentosis, racemis brevibus compositis, bracteis inferioribus herbaceis linearibus racemosis, fructu depresso.

Arbutus tomentosa. *Pursh. fl. amer. sept.* 1. 282. *Hooker & Arnott in Beech. Voy.* 1. 144.

a. hispida; ramis setoso-hispidis tomentosis. *Hooker Fl. Bor. Am.* 2. *t.* 130. *f.* 1. *Bot. Mag. t.* 3320.

β. nuda; ramis tomentosis. *Hooker l. c. f.* 4.

A curious and very rare hardy evergreen shrub, native of rocky places on the west side of North America, from Puget's Sound in the north to California and the Mexican mountains in the south.

It requires to be cultivated in peat and loam, and in a sheltered situation, when it flowers in March; Dr. Hooker informs us that at Glasgow it is kept in the greenhouse. Our specimens were obligingly communicated from the very select collection of William Harrison, Esq., of Cheshunt, where it has been kept in the open air for about four years, and where alone it has as yet flowered in England.

We quite agree with Dr. Hooker in considering the plant which occurs without any bristly hairs upon its branches as a mere variety of this. But we possess two others, like it in habit, which it is important to distinguish. These are

1. *Arctostaphylos cordifolia;* ramis tomentosis, foliis oblongis obtusis subcordatis coriaceis subtus tomentosis supra nitidis, racemis brevibus compositis, bracteis inferioribus foliaceis oblongis coriaceis racemi longitudine, fructu . . .

On the north-west coast of America, Mr. Menzies.

2. *Arctostaphylos glauca;* glabra, glauca, foliis ovato-oblongis acutis coriaceis basi obtusissimis, racemis brevibus compositis, bracteis inferioribus squamæformibus, fructu ovato.

California, Douglas.

There are no plants more general favourites among Collectors than the species of this Natural Order Ericeæ, beginning with Heaths at one end of the series and passing through Andromedas, Vacciniums and Azaleas to Kalmias and Rhododendrons at the other. It is, however, very remarkable, that notwithstanding the extensive commerce of England, the zeal of her merchants, and the enterprize of individuals, some of the most magnificent of these plants are still known to Europeans only from the dried specimens in the herbaria of Botanists. We allude to the noble genus *Befaria*, which contains many species more beautiful than even Rhododendron and Azalea, to the *Thibaudias* with their long tubular crimson blossoms, and to many species of *Gaylussaccia*. The finest of these plants inhabit the Cordilleras of Peru, in the country of the Cinchonas; and certainly if one half the sum that has been sometimes wasted in ill considered undertakings were applied judiciously to an expedition into this region, there would be no reasonable doubt of success, and the results would be indescribably important. We should be glad to explain our views more in detail to any one who might be disposed to engage in such an undertaking, and we cannot help anticipating that the project will be seriously entertained by some of those liberal and wealthy nobles who are indeed becoming the Mæcenates of Natural History in Great Britain. We have been led to these remarks by the examination of a collection of dried specimens lately sent from the Cinchona country by Mr. Mathews, in which we find a larger number of very beautiful plants than we ever have had the good fortune to examine in any collection of similar extent. Among other things there is a most lovely plant which constitutes a new genus related to Thibaudia, and which, as it cannot fail before many years to find its way in a living state to England, we have named in compliment to one of the most liberal of those patrons of science whose noble actions render them the pride of their country as well as the surest pillars of their own high hereditary order. The name of Cavendish has long since been enrolled even in the ranks of practical science, but it shines forth in these later days with renewed lustre in the person of his Grace the present Duke of Devonshire, whose noble residence at Chatsworth is rapidly becoming, by the good taste and munificence of its present proprietor, as remarkable for richness in objects of Botanical importance as it has long been admirable for its extent and princely magnificence.

---

* The English name Bear-berry, or Bear-grape, is a literal translation of the Greek words of which this name is composed.

In order that this plant may be fixed in the records of science we add its technical characters, with a brief description of its general appearance.

CAVENDISHIA.

Calyx superus, campanulatus, truncatus, obsoletè quadridentatus. Corolla tubuloso-cylindracea, 4-dentata. Stamina 8, inclusa, basi imâ corollæ inserta; filamentis carnosis monadelphis; connectivis alternis hastatis auriculis rotundatis, alternis rhomboidalibus. Antheræ muticæ; loculis sulco alto exaratis, apice liberis et rimâ dehiscentibus. Ovarium quadrangulare, 4-loculare, polyspermum. Stigma simplex——*Frutex Peruvianus, sempervirens; gemmis magnis imbricatis; foliis laurinis; floribus speciosis, purpureis, capitatis, involucratis.*

1. Cavendishia nobilis.

This is apparently an evergreen shrub, with broad laurel-like leaves and buds, which are covered over with imbricated scales, very much like those of a Camellia. The flowers are arranged in very close terminal capitate racemes, enclosed within a sort of involucre composed of broad ovate bracts, which are downy on the outside, and coloured red at the edge and in the inside. The corolla is bright crimson and tubular, about an inch long. Sometimes the flowers grow from the side of the old wood, but generally from within involucres at the ends of the branches.

1792

1792



Pub by J. Ridgway 169 Piccadilly Sept. 1. 1835.

S. Watts sc

1792

* CALOTRÓPIS procéra.

Tall Calotropis.

PENTANDRIA *DIGYNIA.*

Nat. Ord. Asclepiadeæ, *R. Br.* (*Introduction to the Natural System of Botany, p.* 210.)

CALOTROPIS.—Suprà, vol. 1. *fol.* 58.

C. *procera;* corollæ laciniis patulis. *R. Br. in Hort. Kew. ed.* 2. 2. 78.
Asclepias procera. *Hort. Kew.* 1. 305. *Willd. sp. pl.* 1. 1263.
Asclepias gigantea. *Andr. Reposit. t.* 271.
Beidelsar. *Prosp. Alp. ægypt. cap.* 25. *c. iv.*

"This plant was raised in the garden of Sir Charles Lemon, Bart., M. P., at Carclew, in June 1832, from seeds collected by Lieut. James Sulivan, R. N., at Porto Praya, St. Jago. In the note which accompanied the seed it is described as being "a shrub or small tree growing from 10 to 20 feet high, and flowering in clusters at the ends of the branches. Leaves six inches long. Pod about the size of the leaf, called by the natives *calmady.*" With us it appears to be a tender, lactescent, upright growing, deciduous shrub, requiring the constant heat of the stove, and thriving in a soil composed of sandy loam, and decayed vegetable earth. It flowered in April.

" *Stem* round, pale green, and together with the leaves thickly covered with hoary pubescence which rubs off on being touched. *Leaves* opposite, decussate, spreading, those of the stem obovate, acuminate, about five inches long and two and a half inches broad. At the extremity of the shoots

---

* Literally "beautifully twisted," apparently in reference to the corolla of C. gigantea.

they are much smaller and more pointed. *Petioles* short, with a deep groove on the upper side. *Flowers* stalked, loosely panicled and spirally arranged, 7, 10, or more in each panicle. *Calyx* 5-parted. *Sepals* very small, stellate, adpressed. *Corolla* divided rather more than half its length, slightly campanulate, and nearly an inch in diameter. When fully expanded it is stellate, and measures about 1¼ inch from point to point of the opposite segments, which are somewhat cordate, acuminate, and exceed a quarter of an inch in breath. Outwardly they are of a pale silvery colour, but inwardly of a deep purplish red, at the point becoming paler and spotted towards the centre of the flower, in which as in others of this Natural Order there is a curious formation, angular at the top, secreting the parts of fructification, and supported by five pale, purplish-tinged, rounded, bracket-like processes. These latter and the divisions of the calyx point in one direction, being placed alternate with the petals and angular part of the centre which point in another. At the extremity of each of the angles is a small aperture, which on being enlarged and having the film removed exhibits two obovate, transparent, greenish yellow bodies, like the pollen masses of Orchideous plants.

"The flowers are produced in succession for several weeks. In the shade, or when the plant is in a room, they are scentless, but in sunshine, or a warm atmosphere, they are highly fragrant."

Mr. Booth has favoured us with the foregoing account and the drawing that illustrates it. The plant is no doubt the *Beid el sar* of Prosper Alpinus, and appears to extend across the whole of the north of Africa, even into Persia, following it may be supposed the desert region till it stops to the westward of Delhi. Alpinus says its juice, which is extremely acrid, was administered successfully as a remedy for ringworm and other cutaneous affections; and that it is also a powerful agent in removing hair from skin.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Oct. 1. 1835.

S. Watts. sc.

1793

* CORYANTHES maculáta.

Spotted Coryanthes.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ, § Vandeæ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

CORYANTHES, Hooker. Perianthium patentissimum. *Sepala* dilatata, flexuosa, conduplicata; lateralibus maximis, basi distinctis. *Petala* multò minora, erecta. *Labellum* unguiculatum, maximum, galeatum, cum basi columnæ continuum, nullo modo articulatum, tridentatum, in medio unguis appendice poculiformi circumdatum. *Columna* teres, basi bicornis, elongata, apice recurva, bialata. *Stigma* rima transversa. *Anthera* bilocularis. *Pollinia* 2, compressa, posticè sulcata, caudiculâ lineari arcuatâ, glandulâ lunatâ apicibus approximato-recurvis.——Herbæ *epiphytæ, pseudo-bulbosæ.* Folia *striata.* Racemi *penduli.* Flores *maximi.* Lindl. Gen. et Sp. Orch. 159.

C. *maculata;* foliis lato-lanceolatis, sepalis lateralibus semicordatis rectis, labello intùs purpureo maculato cornubus elongatis.

C. maculata. *Hooker Bot. Mag. t.* 3102. *Lindl. l. c.*

Sepala *membranacea, tenuia, citò flaccida, supremum lanceolatum acuminatum undulatum, lateralia maxima* (2½-*pollicaria*) *libera, basi angusta, semicordata, primùm explanata vespertilionis alarum more, dein flaccida.* Petala *lanceolata undulata flexuosa in labelli galeam prona.* Labellum *carnosum,* 2½ *poll. longum, unguiculatum, luteo purpureoque variegatum; hypochilio dilatato lateribus inflexis galeato extus rotundato, in mesochilio angusto canaliculato transeunta; epichilio maximo minus carnoso galeato apice ovato inflexo.* Columna *libera, basi cornubus duobus elongatis succum stillantibus instructa, apice clavata et recurva marginibus in alasproductis.* Stigma *rima transversa.*

A native of the woods in Demerara, where it is not uncommon hanging from the branches of trees, and suspending in the air the singular lips of its flowers like fairy buckets, as if for the use of the birds and insects that inhabit the surrounding foliage.

---

* Literally, Helmet-flower; in allusion to the peculiar form of both the hypochilium and epichilium.

There certainly is not a more singular genus than this in the whole vegetable kingdom, nor one whose flowers are less like flowers to the eye of the ordinary observer. The sepals are of the most delicate texture; when young they spread equally round the centre, but after a few hours they collapse, and assume the appearance of a bat's wing half closed. The lip is furnished near its base with a yellow cup, over which hang two horns constantly distilling water into it, and in such abundance as to fill it several times; this cup communicates by a narrow channel formed of the inflated margin of the lip, with the upper end of the latter, and this also is a capacious vessel very much like an old helmet, into which the honey that the cup cannot contain may run over.

C. maculata differs from *C. speciosa*, not only in the colour of the flowers and the greater breadth of its leaves, but also in its lateral sepals being almost exactly half-cordate, without any bend in the middle, and in the horns of the base of its lip being much longer; the middle lobe of its lip too is less distinctly trifid.

C. macrantha, which has lately flowered at Mr. Knight's, surpasses both the others in the richness and magnitude of its flower, and is decidedly distinguished by the mesochilium being irregularly plaited and ridged.

Our drawing of this species was made from a specimen communicated by the Horticultural Society in July 1833.

1794

1794



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Oct 1 1835 J. Watts sc.

1794

* IPOMŒA Aitóni.

Mr. Aiton's Ipomœa.

PENTANDRIA *MONOGYNIA.*

Nat. ord. Convolvulaceæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 218.)

IPOMŒA.—Suprà, vol. 1. *fol.* 9.

I. *Aitoni;* villosa, foliis cordato-subrotundis, trilobis: lobis acutis lateralibus abbreviatis, pedunculis multifloris, bracteis sepalisque divergentibus acuminatissimis, corollâ subcampanulatâ tubo incrassato, staminibus basi glandulosis.

I. Aitoni. *Hort.*

Radix *perennis.* Caulis *villosissimus.* Folia *villoso-velutina; lobo intermedio subrhombeo, acuminato; laminâ petioli longitudine.* Pedunculi *petiolis longiores, multiflori.* Flores *densè aggregati;* bracteis *angustis, canaliculatis, acuminatis, recurvis, villosis, sepalorum longitudine.* Sepala *conformia, corollæ tubo longiora.* Corolla *violacea, lobis brevibus acutiusculis, tubo incrassato sub staminibus glanduloso.* Stamina 5, *ultra tubum exserta, quorum duo longiora.* Ovarium *disco cyathiformi insertum, biloculare; ovulis duobus cuique loculo;* stylus *simplex;* stigma *incrassatum bilobum.*

A pretty perennial stove climber, not uncommon in collections under the name we have adopted. It flowers from April to October. Its native country is unknown to us.

This strictly belongs to the genus Ipomœa, as it has at length been limited by Professor Choisy. The only points in its structure which particularly deserve mention, are the thickened tube of its corolla, and the collection of glands at the base of each stamen, imitating as it were in Ipomœa the scale of the genus Lepistemon.

Increased by seeds and cuttings, the former of which are produced in some abundance.

The flowers open in the morning.

---

* From ιψ a climbing plant, and ομοιος similar; in allusion to the resemblance of this genus to Convolvulus.

1795

1795.



iss Drake. del. Pub by J. Ridgway 169 Piccadilly Oct. 1. 1835. J. Watts sc.

1795

* GOVÉNIA supérba.

Superb Govenia.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

GOVENIA, *Lindl.* *Perianthium* bilabiatum. *Sepala* lateralia falcata, labello supposita, basi paululum connata, supremo paulò majora. *Petala* sub sepalo supremo conniventia, breviora, obliqua. *Labellum* integerrimum, ecalcaratum, concavum, cum basi parum productâ columnæ articulatum, sessile. *Columna* basi paulò producta, teres, subfusiformis, apice utrinque marginata. *Anthera* calyptriformis, 1-locularis. *Pollinia* 4, solida, incumbentia, caudiculâ brevi, glandulâ minore triangulari.——Terrestres. Folia *plicata.* Spicæ *radicales, multifloræ.* Flores *speciosi.* Lindl. gen. et sp. Orch. 153.

G. *superba;* labello ovato-cordato, spicâ cylindraceâ, bracteis acuminatis, foliis oblongis acuminatis basi angustatis scapo æqualibus.

G. superba, *Lindl. in Lodd. Bot. Cab. t.* 1709. *Gen. & Sp. Orch. l. c.*

Maxillaria superba. *Lexarsa et Llave Orch. Mex.* 2. 13.

Caulis *cum foliis 5-pedalis.* Folia 3-*pedalia, et ultra, basi vaginis purpurascentibus vestita.* Racemus 1-1½-*pedalis, cylindraceus, floribus onustus.* Sepala 7 *lineas longa, lateralia semi-oblonga.* Petala *obliquè rhomboidea acuminata.* Labellum 3 *lineas tantum longum, lividum, unguiculatum, cordatum, ovatum, obtusum, suprà canaliculatum.*

This noble species was originally discovered by Messrs. Lexarza and La Llave on the mountains near Valladolid, a town of Mexico, flowering in August. The native name is *Azuzena amarilla.* It is one of the handsomest plants of the whole order of Orchideæ, and is especially remarkable for its stately appearance, the rich orange of its flowers, and the long time they continue to open. The specimen in Messrs. Loddiges' collection grows to the height of a man; that from which our drawing was made was furnished by Mr. Bateman, in February 1834.

---

* So called in compliment to James Robert Gowen, Esq. under whose care were conducted many of the curious experiments upon cross fertilization at Highclere, the seat of the Earl of Carnarvon.

It is found to bear the hottest and dampest stove, but like all the terrestrial species requires a season of repose.

Beautiful as is this plant it must be equalled in the showiness of its blossoms by the *Maxillaria liliacea*, or *Azuzena del Monte* of the people of Valladolid, in the neighbourhood of which town it grows beneath the shade of trees. This plant is described as having a scape about a foot high, terminated by a thryse of snow-white flowers, whose segments are variegated with purple lines, and whose lip is spotted at the base with yellow. It is also in all probability a *Govenia*, and may be named *G. liliacea*.

Of a similar noble aspect is a species collected in Mexico by Count Karwinski, near Atotovilco el Chico, in a cool climate, flowering in June and July. Specimens of this plant, belonging to the Royal Herbarium of Munich, are now before us. The leaves are oblong-lanceolate, tapering to each end, especially to the point, a foot long and about 5 inches wide, and very like those of *Govenia superba*, except that they are smaller. The stem is stated to be five feet high. It is terminated by a short and very dense headed raceme of large flowers, to each of which is an oblong obtuse bract, not quite so long as the ovary. The raceme is 4 inches long, and as much broad at the base. In the form of the flowers it resembles *Govenia superba* very much, but they are three times as large; the sepals are an inch and half long and the other parts in proportion. The lip has an ovate form, but is not cordate at its base, it is not quite half so long as the sepals, and is marked towards its end with largish round very dark dots. The colour of the flowers cannot be ascertained from the dried specimen, but seems to have been pale and uniform; probably pale yellow. This plant may be thus distinguished —

G. *capitata*; labello ovato obtuso, racemo capitato, bracteis oblongis obtusiusculis, foliis oblongis acuminatis basi angustatis scapo multò brevioribus.

1796



[illegible] del.   Pub by J. Ridgway 169 Piccadilly Oct. 1 1835.   [illegible]

* MÍMULUS lúteus; *var.* variegatus.

Variegated Monkey-flower.

DIDYNAMIA *ANGIOSPERMIA.*

Nat. ord. Scrophularineæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 228.)

MIMULUS.—Supra, vol. 11. *fol.* 874.

§. Caule basi sæpius procumbente vel repente, rarius erecto; foliis 3-5-nerviis dentatis; calycibus ovatis tubulosisve sæpiùs inflatis. *Bentham Scroph. Ind. p.* 28.

M. *luteus;* glaber, foliis eroso-dentatis inferioribus longè petiolatis ovatis sublyratisve superioribus rotundatis cordato-amplexicaulibus, calycibus ovatis fructiferis campanulatis dentibus ovatis acutis supremo maximo. *Benth. l c.*

M. luteus. *Linn. Sp. pl.* 884. *aliorumque.*

Var. *variegatus;* corollæ tubo stramineo intus maculato, limbo purpureo.

M. variegatus. *Lodd. Bot. Cab. t.* 1872.

M. luteus variegatus. *Hooker in Bot. Mag.* 3336.

The first plants of this species which appeared in this country were reared from seeds received from Paris; but we believe without any intimation of its native country. Since that time Chile has been stated to be the origin of the species, but we know not upon what certain authority; not a specimen have we ever seen of it from that country. We should almost be inclined to suspect it to be some garden production, if it did not remain true from seed; at all events we agree with Dr. Hooker in referring it to *M. luteus* as a variety.

It is the prettiest of the genus, and is raised with the greatest facility, provided its seed is sown in a damp place under the shade of a wall, or where it is not fully exposed to the sun's rays. So treated it grows vigorously and flowers beautifully in May and June, and will come up again in the same place year after year from its self-sown seeds. But if exposed to too bright a sun, and to dryness, it loses its vigorous habit, and becomes a poor dwindling thing not worth cultivation.

---

* See folio 1330.

It will also strike out freely from cuttings.

1797

1797.



M. Drake del. Pub by J. Ridgway 169 Piccadilly Oct. 1 1835 [illegible]

1797

* PLEUROTHÁLLIS Gróbyi.

The Groby Pleurothallis.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ, §. Malaxideæ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.

PLEUROTHALLIS.—Suprà, vol. 9. *fol.* 759.

P. *Grobyi;* folio obovato emarginato caule triplo longiore racemo laxo erecto multoties breviore, bracteis minimis membranaceis, sepalis costatis oblongis acutis lateralibus apice tantùm sejunctis, petalis lanceolatis acutis, labello lineari obtuso carnoso superne unisulcato.

P. Grobyi. *Bateman in litt.*

Folia *cum caulibus brevibus secundariis cæspitem efficientibus, atroviridia.* Racemi *laxi flexuosi.* Sepala *intus vitellina extus sanguineo vittata.* Petala *flava medio sanguinea.* Labellum *petalis paulò longius, sanguineum apice luteum.* Columna *semiteres, marginata, clinandrio anticè bidentato.* Pollinia 2, *parva, in unum ferè confluentia, basi* (*apice*) *pellucida sed glandulæ nullæ se affigentia.*

A native of Demerara, whence it was imported by Mr. Bateman, who named it in compliment to Lord Grey of Groby, of whose skill and ardour in the cultivation of these curious plants we have already offered our own feeble acknowledgment. (See Grobya Amherstiæ, fol. 1740.)

It is nearly related to *P. picta* from the same country, and with it belongs to a very pretty set of species, all of which are characterized by having their leaves so much longer than the secondary stems as to conceal them and to form dense tufts.

It is readily cultivated in the Orchideous house; but where the atmosphere is not very damp, will require to be kept under a bell-glass.

Our drawing was made at Messrs. Loddiges in April last.

---

* See folio 1298.

We take this opportunity of filling a spare leaf with the characters of some unpublished species of Pleurothallis, together with a new arrangement of the genus, and a few remarks upon *Specklinia* and *Physosiphon*.

PLEUROTHALLIS.

§. 1. *Acaules. Flores aggregati v. solitarii.*

Sp. 1. ruscifolia; 2. laurifolia; 3. succosa; 4. tribuloides; 5. cordata.

6. P. *grandiflora;* folio perfoliato oblongo-lanceolato caudato apice tridentato, pedunculis unifloris fasciculatis, sepalo supremo ventricoso acuminato lateralibus in unum connatis recurvis triplo latiore, petalis lineari-lanceolatis hispido-ciliatis, labello breviore unguiculato subrotundo carnoso ciliato supra unguem excavato.——In *Peruvia;* supra arbores in convalle Lloæ, alt 8000-ped, *Hall.* (*hab. s. sp. comm. cel. Hooker.*)

7. P. *angustifolia;* caulibus ascendentibus folio lineari-lanceolato longioribus, floribus pedicellatis subsolitariis, sepalis carinatis, petalis lanceolatis denticulatis, labello lineari acuto marginibus pone basin incrassatis.——In *Mexico,* prope Xalapam, *Deppe.* (*h. s. sp.*)

8. P. *bidentata;* caulibus ascendentibus folio oblongo-lanceolato apice bi-tridentato longioribus, floribus pedicellatis subsolitariis, sepalis ecarinatis, labello lineari obtuso apice ciliato.——Prope *Rio Janeiro, Forbes.* (*h. s. sp.*)

9. P. *papillosa;* caulibus strictis folio angustè oblongo apice tridentato paulò longioribus, pedunculis subaggregatis unifloris flore vix longioribus, sepalis campanulatis lateralibus semiconnatis supremo oblongo brevioribus, labello subhastato obtusissimo apice vittisq. duabus elevatis carnosis-papillosis——In *Braziliæ* prov. Minarum in arboribus ad S. Joâo Baptista, *Martius* (*ex. s. sp. in hb. M.*) Flores majusculi.

10. P. *discoidea;* caulibus strictis folio ovato-lanceolato brevioribus, pedunculis solitariis unifloris, sepalis oblongis acutis lateralibus omnino connatis, petalis linearibus acuminatis decurvis, labello cordato-ovato undulato obtusiusculo sepalis subæquali.——In insula *Trinitatis* (*v. ic. pict. in hb. Hooker.*) Facies *P. ruscifoliæ.* Flores flavi maculâ sanguineâ oblongâ anticè bidentatâ in medio labelli.

11. P. *punctata;* caulibus strictis folio crasso ovali plano-convexo marginato longioribus, floribus solitariis breviter pedicellatis, petalis lanceolatis apice pubescentibus, labello unguiculato cordato-sagittato obtuso, columna elongata.——In *Brazilia* (*exam. v. c.*). Folia subtus purpureo punctata. Flores atropurpurei. Petala herbacea. Tota planta vix 1¼ poll. alta.

§. 2. *Acaules. Flores racemosi.*

a. *Folia caulibus secundariis longiora* (Cæspitosæ).

12. alpestris; 13? laxa; 14. Grobyi.

15. P. *picta;* folio spathulato marginato retuso racemis laxis duplò breviore, bracteis minimis, sepalis acuminatis lateralibus apice tantùm sejunctis, petalis lineari-lanceolatis acutis, labello lineari obtuso carnoso supra 1-sulcato.——In *Demerara, Loddiges (exam. v. c.).*

16. P. *hymenantha;* folio lineari-lanceolato apice obsoletè 3-dentato racemis flexuosis capillaribus subæquali, bracteis minimis, sepalis acutis lateralibus semisejunctis, petalis acuminatis parum brevioribus, labello oblongo obtuso subundulato complicato.——In *Brazilia;* in arboribus vetustis prope Retiro provinciæ Sebastianop. *Martius. (exam. s. sp. in hb. M.)*

17. P. *ochreata;* folio crasso lanceolato complicato mucrone acuto racemo denso flexuoso nutante duplò longiore, bracteis cucullatis imbricatis acutis, sepalis carinatis acuminatis lateralibus semiconnatis, petalis lineari-oblongis nanis mucronatis, labello . . . . .——In *Brazilia;* in rupibus Serra do Monte Santo provinciæ Bahiensis, *Martius. (exam. s. sp. in hb. M.)* Calyx ruber v. miniatus. Petala duplo minora flava, omnia subæquali. *Mart. mss.*

18. P. *rupestris;* folio tereti mucronato anticè sulcato racemo stricto rigido paulò breviore, bracteis parvis rigidis concavis, sepalis acutis lateralibus semiconnatis, petalis lanceolatis ciliatis, labello subtrilobo membranaceo obtuso appendice magno carnoso bifurco: cruribus parallelis porrectis acutis.——In *Brazilia,* inter rupes M. Itacolumi et Moro de Vª. Rª. provinciæ Minarum, *Martius. (exam. s. sp. in hb. M.)* Scapus rubenti-virens. Sepala purpurascentia. Petala purpurea. *Mart. mss.*

19. P. *teres;* folio tereti racemo gracili breviore, bracteis membranaceis ochreatis, floribus pendulis, sepalis acuminatis lateralibus apice tantum liberis, petalis minimis oblongis acutis, labello lineari canaliculato acuto marginibus incrassatis, columnâ basi pubescente.——In *Brazilia, Loddiges (exam. v. c.).* Flores cinnamonei. Folia omnino *Leptotis bicoloris.*

20. P. *hians;* caule brevi, folio oblongo basi angustato obtuso racemo flexuoso 6-9-floro duplò breviore, bracteis minutis ochreatis, floribus bilabiatis: labio postico 2-dentato antico oblongo obtuso, petalis cuneatis, labello ovato acuto complicato.——In *Brazilia,* in montibus Organ *(v. ic. pict. et exam. fl. s. in herb. Hooker.)* Caulis pollicaris. Folium 3-pollicare. Scapus fusco purpureus. Flores ejusdem coloris, aciculatìm maculati, extùs pallidè cinnamonei. Sepalum supremum anticum, lateralibus connatis posticis æquilatum. Petala labellumq. nana atropurpurea. Columna ejusdem coloris clinandrii margine membranaceo latissimo.

21. spiralis. 22. racemiflora.

b. *Folia caulibus secundariis breviora v. æqualia.* (Macropodæ.)

23. prolifera; 24. saurocephala; 25. quadrifida (=Dendrobium quadrifidum. *La Llave*); 26. pulchella; 27. Lanceana.

28. P. *Jamiesoni;* folio lineari-oblongo carnoso submucronato racemo secundo stricto duplò breviore, bracteis densis brevibus ochreatis cucullatis, sepalis

oblongis subæqualibus leviter carinatis lateralibus basi tantum connatis, petalis duplo brevioribus apice rotundatis, labello breviore membranaceo rhombeo-trilobo plicâ unguis transversâ.——In *Peruvia*, in præcipitiis Pichinchæ haud procul a Quito, *Jamieson*. (*exam. s. sp. in hb. Hooker.*) P. pulchellæ valdè similis, sed minor.

29. P. *capillaris;* folio lineari-lanceolato racemis capillaribus fasciculatis subæquali, sepalis acuminatis lateralibus basi connatis, petalis ferè æquilongis serratis acuminatis, labello lanceolato basi angustato acuto membranaceo 3-venio. ——In *Brazilia*, in arborum truncis provinciæ Sebastianop. *Martius*. (*exam. s. sp. in hb. M.*) Facies P. *ruscifoliæ*.

30. lanceolata.

31. P. *sclerophylla;* folio oblongo petiolato obtuso carnoso racemis multò breviore, bracteis ovatis cucullatis, sepalis acuminatissimis omnibus ferè liberis, petalis nanis obtusis, labello oblongo conduplicato apice piloso-glanduloso.—— In *Brazilia;* in Serra de Piedade, prov. Minarum, etiam in rupibus campestribus prov. S. Pauli, *Martius*. (*exam. s. sp. in hb. M.*)

32. P. *rubens;* folio oblongo-lanceolato obtuso, racemis longissimis, bracteis oblongis ochreatis, sepalis lanceolatis acuminatis lateralibus basi tantùm connatis petalis obtusis, labello obovato repando obtuso lineis 2 elevatis flexuosis carnosis. ——In *Brazilia;* in rupibus ad Opp. S. Joao d' El Rey, *Martius*. (*exam. s. sp. in hb. M.*)

§. 3. *Caulescentes;* i. e. rhizomate nullo sed caule folioso.

33. macrorhiza. 34. caulescens.

Under the name of *Stelis tubata*, Messrs. Loddiges have published a very curious plant with the habit of Pleurothallis, but with its sepals all united into a tube, which is inflated at the base and contracted at the mouth. Otherwise its fructification has the structure of *Stelis*. I propose to distinguish this as a peculiar genus, with the following name and character.

PHYSOSIPHON.

Calyx tubulosus basi ventricosus, apice 3-fidus. Petala in fundo calycis carnosa, nana. Labellum et columna Stelidis. Pollinia 2, sphærica.——Herbæ epiphytæ habitu Pleurothallidis.

1. P. Loddigesii; (=Stelis tubata, *Lodd. Bot. Cab. t.* 1601.)

2. P. emarginata; (=Pleurothallis emarginata, *Lindl. Gen. et. Sp. Orch. Pl. p.* 6.)

3. P. *spiralis;* folio angustè oblongo subemarginato carnoso spicâ spirali confertâ duplò breviore, bracteis minimis ovatis acutis cucullatis, calyce semitripartito laciniis erectis acutis, petalis cuneatis carnosis, labello ovato-subrotundo membranaceo apiculato trivenio.——In *Brazilia*, ad Portum S^{æ}. Catharinæ, *Tweedie*. (*hab. s. sp. comm. Cel. Hooker.*)

SPECKLINIA.

To this genus is to be added the following:—

S. *atropurpurea;* vaginis caulis 2-3 ventricosis, folio oblongo plano 7-9-venio, flore solitario subsessili, petalis 3-cuspidatis, labelli laminâ sagittatâ.——In *Jamaica.* (*Exam. s. cult. in herb. Hooker. ex horto Liverp.*) Folium 2½ poll. longum, basi angustatum. Vaginæ infundibulares, pollicem longæ, in caulem 2-pollicarem. Flores atropurpurei. Ovarium lineam longum, hexapterum; alis apice truncatis petalinis minoribus. Alabastrus quem tantum vidi semunciam longus, angustus, falcatus. Sepala anteriora basi gibbosa, oblongo-linearia. Petala nana, vix 2 lineas longa, ovata, 3-cuspidata, obliqua, sæpiùs venâ alterâ excurrente, alterâ in dimidio majore petali abbreviatâ. Labellum unguiculatum, in columnam pronum, cum ejus basi articulatum; ungue lineari, 3-venio axi cristato, laminâ sagittatâ apicibus posticis obtusis. Columna petaloideo-alata, apice tridentata, basi paulò producta; pedis marginibus carnosis rotundatis medio unidentatis. Anthera parva, 1-locularis. Pollinia 2, pyriformia, atropurpurea, apice materie diaphanâ granulosâ stigmati adhærentia.

And probably several spurious Dendrobia of authors, especially—

S. retusa = Dendr. retusum, *La Llave.*
S. scariosa = Dendr. scariosum, *Do.*
S. pusilla = Dendr. pusillum, *Humb. et Kunth.*
S. acuminata = Dendr. acuminatum, *Do.*
S. elegans = Dendr. elegans, *Do.*

1798

1798.



Miss Drake del.

Pub by J Ridgway 169 Piccadilly Oct. 1. 1835.

S. Watts sc.

* EDWÁRDSIA chilénsis.

Chilian Mayu Tree.

DECANDRIA *MONOGYNIA.*

Nat. ord. LEGUMINOSÆ, *Juss.* § PAPILIONACEÆ. (*Introduction to the Natural System of Botany, p.* 87.)

EDWARDSIA.—Suprà, vol. 9. *fol.* 738.

E. *chilensis;* foliolis 13-19 elliptico-oblongis obtusis rigidis subtus sericeis, petalo supremo lateralium longitudine, legumine binodoso aptero.
Edwardsia chilensis. *Miers Trav. in Chili; the name only.*
Sophora macrocarpa. *Smith in Rees.* 7t/9 *De Cand. Prodr.* 2. 96. *Hooker et Arnott in Bot. Misc.* 3. 177.

A fine tree, native of Chile, where the inhabitants call it *Mayu.* It was introduced by Messrs. Loddiges in 1822, having been sent them by General Paroissien under the name of "*Sophora Myrospermum.*"

It proves to be a hardy handsome tree, flowering in May in great profusion. We believe that it at present exists in no other collection.

This is plainly an *Edwardsia,* notwithstanding the opinion of the late Sir James Smith to the contrary, and notwithstanding the alleged want of wings to its pods. Supposing that fact to be as stated, it only shews that the winged pod is unessential to the genus Edwardsia, and not that the genus itself is a bad one; for surely if there be any one genus on which Nature has set her mark more distinctly than another it is this. Mistakes like those of Smith are the certain accompaniments of the meagre system of artificial Botany.

---

* See folio 738.

It is the short loose axillary racemes, the hard cup-shaped calyx, slit on the upper side, and the broad vexillum which overlaps the other petals and is usually absolutely parallel with them, that distinguish *Edwardsia* from *Sophora*, whose racemes are terminal, calyx soft and regularly toothed, and vexillum narrow and recurved, or reflexed.

1799

1799.



S. Drake del.   Pub by J. Ridgway 169 Piccadilly [illegible]   J. Watts sc.

* MAXILLÁRIA crócea.

Saffron-coloured Maxillaria.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

MAXILLARIA.—Suprà, vol. 11. *fol.* 897.

§. 2. *Scapigeræ.* Pedunculi radicales. Labellum anticum. Scapi uniflori.

M. *crocea;* pseudobulbis oblongis compressis foliatis, foliis oblongis undulatis obtusis emarginatis latè vaginantibus, scapis erectis unifloris laxè vaginatis foliis duplo brevioribus, sepalis petalisque angustè triangularibus apice elongatis abruptè acutis, labello obovato obtuso levissimè trilobo apice carnoso crispo supra medium unituberculato.

We have received the drawing and following account of this plant from Mr. Booth.

"We owe the introduction of this curious plant to Capt. Sutton, of his Majesty's Packet establishment at Falmouth, who imported it from Rio Janeiro in the spring of 1833, and presented it, with others, to Sir C. Lemon, Bart., M. P., in whose collection at Carclew, Cornwall, the subject of the present notice produced its flowers in August 1834.

"*Root* fibrous. *Pseudo-bulbs* small, oblong, compressed, nearly covered by the three sheathing, strap-shaped leaves, which are each thick and rigid, from four to six inches long, and an inch in width, a little twisted, with an oblique sharp point. The leaf on the bulb is similar to the others but much longer. All of them are of a deep green above, and a pale silvery green beneath. *Scapes* four inches high, one-flowered, round and slender, rising from the base of the leaves, and covered with three or four, thin, pale green, acuminate bracts, forming a joint at their junction with the stem. *Flowers* before expansion having some resemblance to the beak of a bird, pale green with a brownish tinge at the spur. *Perianth* of a greenish yellow, becoming of an orange colour when expanded. *Sepals* narrow and pointed, the three exterior ones being each about the same size and length, with their edges turned back. The two interior ones are rather shorter than the others, somewhat curved, and crossing so as almost to conceal the column. *Labellum* about half the length of the sepals, thick and fleshy, connected to the lengthened base of the column, which forms the spur; cucullate and crisp at the margin, which is rather paler than the sepals, otherwise it is of a brownish red colour, together with the column.

---

* See folio 1428.

"It is obviously allied to *M. picta* and *punctata*, from both which the character assigned to it, abundantly distinguish it.

"Like most of the terrestrial Orchideous plants from Tropical countries, it requires the constant heat of the stove, and to be kept rather moist. It seems to flourish in the soil we find in the hollow decayed trunks of trees, mixed with some finely chopped moss; and promises to increase freely by off-sets."

1800.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Oct 1, 1835.

J. Watts sc.

* STANHÓPEA oculáta.

Eyed Stanhopea.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

STANHOPEA.—Suprà, vol. 18. *fol.* 1529.

S. *oculata ;* labello medio constricto, hypochilii elongati lateribus cornutis erectis basi bimaculatis, epichilio ovato acuminato cornubus duobus baseos incurvis. *Lindl. gen. et sp. Orch. Pl. p.* 158.

Ceratochilus oculatus. *Lodd. Bot. Cab. t.* 1764.

Imported by Messrs. Loddiges. It is the most interesting of this splendid genus, on account of the extremely delicate waxy appearance of its surface, the softness of its ground colour, and the richness of the deep purple spots, which lying upon a bright yellow field so very conspicuously ornament the base of the lip and the petals.

Our drawing was taken from a most noble specimen produced in July 1834, in the stove of Mr. Bateman. The leaves, including the petioles, were more than two feet long, and being of the darkest green had a most imposing appearance. Mr. Bateman informs us that he finds it requires eight such leaves to form a flowering plant; and that those who wish to succeed in blossoming it must suffer it to remain a long time undisturbed, since a good many pseudo-bulbs are required to furnish one really fine specimen.

The singular form of the labellum of this and some other Orchideous plants has rendered it necessary to distinguish it into three parts for the convenience of description.

---

* See folio 1529.

The lowest part of the lip is the *hypochilium*, the upper end the *epichilium*, and the intermediate portion the *mesochilium*. All these parts are, however, nothing more than divisions in a very fleshy lip, produced by contractions of its sides, and by the lobes, so common in the order, variously arranged and combined.

The species is at once known by the narrowness and length of the hypochilium, independently of all other circumstances.

It is said to be a native of Brazil; but we suspect some mistake in this statement. It is certainly wild in Mexico, for it exists among Count Karwinski's plants in the Royal Munich Herbarium.

1801

1801.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Oct. 1. 1835. J. Watts sc.

* PRÚNUS japonica.

The Single Chinese Plum.

ICOSANDRIA *MONOGYNIA.*

Nat. ord. Rosaceæ, § Amygdaleæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 81.)

PRUNUS.—Suprà, vol. 2. *fol.* 136.

P. japonica. *Supra, vol.* 1. *fol.* 27. The double state.

It is always interesting to procure the wild forms of cultivated species, and so see from what humble originals Nature produces some of the most striking of her works.

The double Chinese Plum, or Almond, as it is often incorrectly called, is beyond all comparison the handsomest plant of its season; there is nothing to vie with its crowded clusters of most delicate blushing flowers, the petals of which are loosely, but symmetrically, arranged into the most perfect of vegetable beauties. The simple shrub now represented is its origin, and is one of the many examples of the creation by the patient Chinese of the fairest ornaments of the garden, from the most inconspicuous plants of the woods.

For its introduction the public is indebted to John Reeves, Esq., by whom so large a proportion of all the fine Chinese flowers now common all over Europe have been procured for this country.

It appears to be a hardy shrub; our specimen, however, was taken in January last, from a plant which had flowered in a greenhouse.

The double state of the species is represented in the first volume of this work.

---

* See folio 1243.

1802

1802.



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Oct. 1 1835. J. Watts sc.

* MAXILLÁRIA picta.

Painted Maxillaria.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

MAXILLARIA.—Suprà, vol. 11. *fol.* 897.

---

§. 2. *Scapigeræ.* Pedunculi radicales. Labellum anticum. Scapi uniflori.
M. *picta;* pseudobulbis ovatis subsulcatis 1-2-phyllis, foliis oblongo-lanceolatis planis, scapis erectis unifloris distanter vaginatis foliis duplò brevioribus, sepalis petalisque lanceolatis (abruptè acutis) sub-conniventibus, labello trilobo medio unituberculato lobis oblongis obtusis lateralibus brevioribus.
M. picta. *Hooker in Bot. Mag. t.* 3154. *Lindl. gen. & sp. Orch. p.* 146.

A truly beautiful plant found wild in the Organ mountains of Brazil, and originally introduced by the late lamented Mrs. Arnold Harrison. When well managed it throws up a profusion of its gay speckled flowers, which remain perfect for some time.

Our drawing was made in the stove of the Messrs. Loddiges in November 1834.

There are two other unpublished species of Maxillaria, belonging to this section, which we take this opportunity of naming.

1. M. *rufescens;* pseudobulbis ovatis subtetragonis monophyllis, foliis lanceolatis utrinque acuminatis, scapis unifloris prostratis vaginis minimis distantibus, sepalis petalisque oblongis conformibus obtusis, labello oblongo trilobo etuberculato laciniis lateralibus minimis acutis intermediâ elongatâ emarginatâ.

A native of Trinidad, whence it was imported by Mr. Lowe of Clapton. Our description is taken from a drawing and flower transmitted from Chatsworth in December 1834.

---

* See folio 1428.

It is a species of no beauty. The sepals are a dull greenish purple, the petals and lip yellow, the latter speckled with dull purplish red. Possibly the specimen was bleached, and the colours would become brighter if the plant had flowered in brighter weather. Stands next M. *picta*.

2. M. *graminea*; pseudobulbis nullis, foliis gramineis recurvis, perianthio campanulato, sepalis oblongis obtusis lateralibus paululum connatis, petalis angustioribus, labello cum pede longè producto columnæ articulato trilobo: lobis lateralibus semi-ovatis acutis ascendentibus intermedio majore reniformi, tuberculo disci magno carnoso truncato posticè bilobo.

A native of Demerara; imported by Mr. Lowe; flowered in Messrs. Loddiges' collection, January 1835. A small and inconspicuous species. The blossoms are pale yellow, with the segments a little banded with purple near the base. Not very nearly related to any known species.

1803



Miss Drake del.   Pub. by J. Ridgway 169 Piccadilly Oct. 1. 1835.   J. Watts sc.

1803

*CHIRÓNIA peduncuLáris.

Long-stalked Chironia.

PENTANDRIA *MONOGYNIA.*

Nat. Ord. Gentianeæ, *Juss.* (*Introduction to the Natural System of Botany, p.* 215.)

CHIRONIA.—Suprà, vol. 3. *fol.* 197.

C. *peduncularis;* glabra, foliis ovato-lanceolatis acuminatis sessilibus 3-5-veniis internodiis subbrevioribus, pedunculis unifloris foliis longioribus, calyce tubo corollæ breviore, caule tereti.

C. trinervis. *Hort. nec Linn.*

Frutex *diffusus,* 3-4-*pedalis, atroviridis, glaber, succo amarissimo sublacteo scatens.* Internodia *basi purpurascentia.* Folia 3-5-*venia,* 2-*poll. longa, sæpe internodiis breviora.* Pedunculi *axillares et terminales, foliis duplò longiores, uniflori.* Calyx 5-*partitus, laciniis linearibus acuminatis, tubo corollæ brevioribus.* Corolla *persistens, tubo gracili viridi semunciali, limbo patente* 5-*partito, purpureo, ejusdem longitudinis; laciniis ovatis acutis post anthesin convolutis.* Stamina 5, *fauce tubi inserta, a stylo aversa, exserta;* antheris *linearibus poro duplici apicis dehiscentibus, defloratis rectis.* Ovarium *teres, acuminatum, uniloculare marginibus carpellorum* 4 *inflexis ovuliferis;* stylo *continuo;* stigmate *dilatato bilobo.* Fructus *semimaturus carnosus, amarissimus.*

A very pretty greenhouse shrub, covered with a succession of purple flowers from July to October, and propagated freely by cuttings.

Nothing can be easier than its management, as it grows in any kind of soil, will thrive out of doors in summer, and will survive the winter without injury in a very indifferent greenhouse.

---

* Named in honour of Chiron the Centaur, one of the earliest medical practitioners whose names have been preserved. The intense bitterness of the genus indicates its powerful tonic properties.

We are unacquainted with its history; it does not seem to have been before described; and we have not been able to learn even its native country. In gardens it is called Chironia trinervis, but it is a totally different plant from the Ceylon species so named by Linnæus, which appears to be a genuine Exacum, as that genus is limited by Dr. Brown in his Prodromus.

The bitterness of the species is most remarkable, even mong its bitter neighbours.

1804

1804



Miss Drake del.   Pub. by J. Ridgway 169 Piccadilly Oct 1 1835.   J. Watts sc.

* MAXILLARIA densa.

Dense-flowered Maxillaria.

◆

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

MAXILLARIA.—Suprà, vol. 11. *fol.* 897.

---

§. 1. *Axilliflorœ.* Pedunculi axillares.

M. *densa;* pseudobulbis oblongis compressis axillaribus monophyllis, foliis oblongo-lanceolatis obtusis emarginatis, racemis axillaribus densissimè aggregatis, bracteis cucullatis, perianthiis bilabiatis, sepalis lineari-lanceolatis acuminatis carinatis, petalis paulo minoribus, labello oblongo indiviso apice recurvo et canaliculato, medio lineâ transversâ elevatâ.

A native of Mexico, whence it was imported by the Messrs. Loddiges, in whose stove it flowered in January last.

It belongs to that set of Maxillarias on which the genus was founded by the authors of the Flora Peruviana, and which in some respects appear to differ from those with which we are but acquainted in Gardens. The stems rise some height above the ground, and are closely invested with brown withered scales, from the axils of which spring the pseudo-bulbs and flowers; in most Maxillarias on the contrary, there seems to be no stem, but the pseudo-bulbs sit close upon the ground, as in *M. picta* and *ochroleuca.* In truth, however, there is a stem in both kinds; only it is erect and above ground in the one, and prostrate or underground in the other. And so it is with all the pseudo-bulbous Orchideæ; their true stem is in most cases a rhizoma, like that of Iris, rooting on the side that touches the ground, and pushing up leaves and pseudo-bulbs on the other.

---

* See folio 1428.

1805



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Oct. 1. 1835. S. Watts sc.

1805

*FÚCHSIA díscolor.

Port Famine Fuchsia.

◆

OCTANDRIA *MONOGYNIA.*

Nat. ord. ONAGRACEÆ, *Juss.* (*Introduction to the Natural System of Botany, p.* 56.)

FUCHSIA.—Suprà, vol. 10. *fol.* 847.

F. *discolor;* dumosa, ramosissima, vinosa, ramulis levissimè pubescentibus, foliis ovatis denticulatis undulatis petiolatis oppositis ternatisque, pedunculis foliis longioribus, petalis obtusis convolutis calycis laciniis acuminatis brevioribus, staminibus longè exsertis.

A native of Port Famine in the Falkland Islands, whence it was some years since introduced by Mr. Lowe. Our drawing was made from a plant in the Garden of the Horticultural Society in August last. It has the same season of flowering as the other Fuchsias, and may be increased readily by cuttings in like manner.

If we are asked to state in what respects this differs botanically from *F. gracilis* and *tenella*, we should find it very difficult to answer the question. And yet it is a decidedly different plant; remarkable for its compact, bushy manner of growth, its deep purple branches, its small very undulated leaves, and also for its being apparently more hardy than any other Fuchsia yet in the Gardens. For the latter reason we attach especial importance to it, for by a judicious intermixture of its pollen with such beautiful plants as *F. conica, globosa,* and its other more tender relatives, the whole race may probably be rendered capable of bearing the climate of Great Britain, and may thus become far more generally valuable than they yet are.

---

* See folio 1269.

It is probable that those are right who consider the greater part of the Chilian Fuchsias mere varieties of *F. macrostema*, and if so this will have to be added to the list as a dwarf contracted kind. *F. globosa* and *conica* are, however, to all appearance distinct species.

1806



1

2

M. Drake del.

Pub by J. Ridgway 169 Piccadilly Oct 1 1835.

J. Watts sc.

1806

FERNANDÉZIA acúta.

* *Sharp-leaved Fernandezia.*

◆

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchideæ, §. Vandeæ, *Lindl.* (*Introduction to the Natural System of Botany.* p. 262.)

FERNANDEZIA, Fl. Peruv. Perianthium patens. *Sepala* libera. *Petala* conformia, sub sepalo supremo conniventia. *Labellum* anticum v. posticum, ecalcaratum, liberum, trilobum, disco tuberculatum. *Columna* brevis, utrinque aurita. *Anthera* bilocularis. *Pollinia* 2, solida, obovata, *glandulâ* . . . . . *caudiculâ* . . . . . .——Epiphytæ *caulescentes.* Folia *disticha, equitantia, imbricata.* Racemi *pauciflori, terminales v. laterales.* Flores *parvi lutei.* Lindl. Gen. et Sp. Orch. p. 207.

F. *acuta;* foliis acuminatis carinatis, corymbo laxo multifloro, bracteis obtusis membranaceis, labello lineari 3-lobo laciniis lateralibus rectis abbreviatis intermediâ emarginatâ obtusâ, disco pulvinato tuberculo minimo pone basin.

Omnia fere F. elegantis *nisi folia et labellum. Hoc pallidè luteum disco prominulo pulvinato anticè bilobo sanguineo-marginato; laciniis lateralibus nullo modo hastæ effigiem referentibus sed laciniâ intermediâ parallelis.*

A native of Trinidad, whence it was imported by Mr. Knight, in whose collection it flowered in June last.

In habit is very like *F. elegans,* from which it is principally distinguished by the tapering form of its leaves, and the shape of its lip, which, instead of having a sort of hastate form, in consequence of the divergence of its lateral lobes, and a number of irregular unequal tubercles upon its disk, has the lateral lobes parallel with the middle one, only a single small tubercle at its base, and a slightly elevated disk which is 2-lobed, and bordered with crimson in front.

A hot damp stove is necessary to keep this rare species in health.

---

* So called after George Garcias Fernandez, a Spanish Botanist, who is unknown, except through the medium of this genus.

1807

1807.



Miss Drake. del.

1807

* CÉREUS triangúlarís.

Triangular Torch-thistle.

ICOSANDRIA *MONOGYNIA.*

Nat. ord. CACTACEÆ. (*Introduction to the Natural System of Botany, p.* 54.)

CEREUS, D. C. *Sepala* numerosissima, imbricata, basi ovario adnata, in tubum elongatum concreta, exteriora breviora calycinalia, media longiora colorata, intima petaliformia. *Stamina* numerosissima cum tubo concreta. *Stylus* filiformis apice multifidus. *Bacca* sepalorum reliquiis areolata, tuberculata aut squamata. *Cotyledones* nullæ? (depauperatæ).——Frutices *carnosi, elongati, axi ligneo internè medullifero donati, angulis verticalibus spinarum fasciculos gerentibus regulariter sulcati.* Anguli *seu* alæ *nunc plurimæ, nunc paucissimæ, rariùs duæ tantum et tunc rami compresso-alati.* Flores *ampli è spinarum fasciculis aut crenis angulorum orti.* De Cand. Prodr. 3. 463.

§. 2. *Serpentini.* Caulis articulatus prostratus radicans vel volubilis. D. C.

* * * 3-4-*angulares.*

C. *triangularis ;* repens trigonus, aculeis brevibus quaternis decussatis, sepalis exterioribus spatulatis apice foliaceis, petalis lanceolatis cuspidatis.
Cactus triangularis. *Linn. sp. pl.* 666. *Jacq. amer.* 152.
Cereus triangularis. *Haworth. syn.* 180. *De Cand. prodr. l. c.*

A native of Mexico and the West India Islands, whence it was introduced long since, but it flowers so rarely that it has never yet been represented from an European specimen. Bradley, who has given it in his work on succulent plants, has only figured its stem.

For the specimen from which our drawing was made we are indebted to Sir George Staunton, Bart., in whose garden at Leigh Park, near Havant, it flowered in September 1834, under the good management of Mr. Robert S. Wilson the gardener. The plant had been in the collection at Leigh Park upwards of fifteen years without blossoming; between March and September it produced shoots upwards of seven feet long.

---

* Literally, a torch or taper; a name translated by the English Torch-thistle; and given to these plants in consequence of the upright kinds having something the appearance of the tapers used in the ceremonies of the Roman Catholic religion.

Two flowers were perfected, of which one opened about six o'clock in the afternoon of the 22nd of September, and faded about eleven A. M. on the following day; the other was despatched to London by coach on the 24th of the same month, and reached London in perfect condition.

It was indeed a beautiful object, its petals were of the most dazzling whiteness, the effect of which was greatly heightened by the dense mass of yellow stamens occupying the centre, and by the border of olive green sepals, on which the petals reposed. This is said to have the largest flowers of all the species, not even excepting the common night-blowing Cereus; its fruit is described as being quite smooth, of a rich scarlet, and with the size and form of a goose's egg.

1808

1808.



Miss Drake del.

Pub. by J. Ridgway 169 Piccadilly Nov. 1 1835.

Watts sc.

1808

* EÚTOCA víscida.

Clammy Eutoca.

PENTANDRIA *MONOGYNIA.*

Nat. ord. HYDROPHYLLACEÆ. (*Introduction to the Natural System of Botany. p.* 244.)
EUTOCA.—Suprà, vol. 14. *fol.* 1180.

E. *viscida ;* glanduloso-pilosa viscosa, caule erecto ramoso, foliis cordato-ovatis subangulatis serratis, racemis elongatis furcatis simplicibusque, placentis multi-ovulatis.
E. viscida. *Bentham MSS.*

Annua, undique pilis nigro-capitatis glandulosis viscosa; caule tereti ramoso bipedali. Folia *tactu mollia et oleosa, 2 p. lata, paulò longiora, sensim decrescentia, superiora grosse simpliciter dentata.* Racemi *gyratim elongantes, multiflori, ebracteati.* Sepala *linearia, obtusa, capsulæ longitudine.* Corolla *generis, amœne cœrulea tubo roseo.* Filamenta *setacea basi villosa ;* antheræ *flavæ.* Capsula *ovata, mucronata, apice bivalvis, semibilocularis, placentis parietalibus polyspermis.* Semina *minuta, fusca, scrobiculata, marginibus areolarum acutis centro foveola excavata.*

A new hardy annual, found in California by Mr. Douglas, and raised by the Horticultural Society, in whose garden it flowered this summer for the first time. It is perfectly hardy and will grow in any common soil. Our drawing was made last July.

The blue of the flowers is remarkably deep and brilliant, so that the plant has a handsome appearance ; but its leaves are rather coarse and weedy. We know, however, of no plant better adapted for bouquets ; for it will go on growing and flowering in water for two or three weeks after being gathered.

The whole of its surface is covered with hairs, having little black heads filled with a viscid secretion, and sticking to the fingers like those of the Henbane ; these are so crowded about the pedicels as to give them quite a sooty aspect.

---

* See folio 1180.

1809

1809.



S. Watts del. Pub by J. Ridgway 169 Piccadilly Nov. 1. 1835. J. Watts sc.

* VÁNDA téres.

Taper-leaved Vanda.

GYNANDRIA *MONANDRIA.*

Nat. ord. Orchidaceæ, § Vandeæ. (*Introduction to the Natural System of Botany, p.* 262.)

VANDA.—Suprà, vol. 6. *fol.* 506.

V. *teres;* foliis teretibus, racemis ascendentibus subbifloris foliis æqualibus, sepalis oblongis obtusis: supremo erecto lateralibus semitortis labello suppositis, petalis majoribus suborbiculatis undulatis, labello basi conico: laciniis lateralibus ascendentibus subtruncatis intermediâ pubescente apice dilatatâ truncatâ emarginatâ. *Lindl. gen. et sp. Orch. p.* 217.

This superb epiphyte was originally discovered upon trees in Sylhet by Dr. Wallich, and it has more recently been met with by Mr. William Griffith abundantly near Medown in the Burmese Empire, also growing on trees in the woods.

When Dr. Wallich came to England, this species formed one of the numerous living plants brought home under his care, and blossomed during the voyage; the very flowers which were then produced we are so fortunate as to possess in spirits, through the kindness of their liberal discoverer; they measure nearly four inches and a half from the tip of one petal to that of the other, but only two were formed upon the spike. The specimens from Mr. Griffith are not quite so large in the flower, but there are three blossoms to each spike.

---

* Vandá is the Sanscrit name of the original species of this genus.

That from which our drawing was taken was produced at Syon, in the collection of his Grace the Duke of Northumberland, by whose permission we are enabled to publish it. It was not quite so fine as what we have represented, but considering what the native specimens are we have felt quite justified in forming our figure between the two. The flowers in the accompanying plate, though larger than those on the plant at Syon, are considerably smaller than those we possess in spirits.

Nothing can exceed the flowers of this plant in delicacy of texture or softness of colour, the deep purple of the petals softens away to the margin, and seems to melt as it were into the purer white of the sepals, while the rich crimson and yellow of the lip renders the brilliancy of the other parts still more conspicuous.

We found the plant had been managed by having its stem covered with moss, which if kept damp would maintain the surface in a uniform state. We do not, however, perceive any trace of the roots in our wild specimens having been growing among moss, and we consider it probable that it will succeed as well if the roots are exposed freely to the air.

1810



S. Watts sc. Pub by J. Ridgway 169 Piccadilly [illegible] 1835 S. Watts sc.

*CRATǼGUS Douglȃsii.

The Douglas Thorn.

ICOSANDRIA *MONO-PENTAGYNIA.*

Nat. ord. Rosaceæ, § Pomeæ. (*Introduction to the Natural System of Botany, p.* 83.)

CRATÆGUS.—Suprà, vol. 83. *fol.* 1128.

C. *Douglasii;* foliis obovatis ovalibusque inciso-serratis acutis basi cuneatis glabris demum subcoriaceis sublucidis, laciniis calycis ovatis glanduloso-serratis pedunculisque glabris, spinis rigidis rectiusculis.

C. punctata. *β.* brevispina. *Douglas in Hook. Fl. Bor. Am.* 1. 201.

Rami *ascendentes, cinereo-purpurei, glabri, spinis rigidis lævigatis, nunc brevibus nunc longissimis.* Folia *semper glabra, petiolis omninò eglandulosis, inciso-serrata nullo modo angulata vel plicata, sub fine anni coriacea et sublucida, citò tamen decidua, paulò post* C. glandulosam *et* punctatam. Inflorescentia *cymosa glabra,* floribus *magnitudine mediocri subtetragynis.* Sepala *ovata glanduloso-serrata.* Fructus *parvus, atropurpureus.*

A native of North West America, where it was collected by Mr. Douglas. A hardy tree of small size flowering in May.

We believe this to be essentially different from all the published species of this genus, but the whole of the synonymy is in so deplorable a state of confusion, from want of some monographer, who at the same time is well acquainted with the living trees, and has access to authentic sources of information, that we are by no means certain that it has not already appeared in print as a species.

Mr. Douglas considered it a variety of C. *punctata,* but the leaves of that species are far thinner, and have an evident tendency to be furrowed as if plaited; its branches are horizontal, and its fruit large and either red or yellow, in all which respects it differs from this. Its inflorescence, moreover, is downy. It is possible that what is called C. *macracantha* in the gardens may not be specifically distinct, but as its fruit is red it requires further examination.

---

* See folio 1161.

The leaves of this species are remarkably leathery in the autumn, and then acquire a purplish cast and are shining. They fall off about the same time as those of C. *punctata* and *pyrifolia*.

The plant from which our drawing was taken stands in the Arboretum of the Horticultural Society.

1811

1811.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Nov. 1. 1835. J Watts sc.

* MAXILLÁRIA cristâta.

Crested Maxillaria.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDACEÆ, § VANDEÆ. (*Introduction to the Natural System of Botany, p.* 262.)

MAXILLARIA.—Suprà, vol. 11. *fol.* 897.

M. *cristata;* pseudobulbis ovatis sulcatis monophyllis, foliis oblongo-lanceolatis plicatis, scapo pendulo bifloro squamis laxiusculis vaginato, floribus explanatis, sepalis petalisque lanceolatis acutis æqualibus, labello multo minore tripartito carnoso: laciniis lateralibus falcatis intermediâ rotundatâ cristato-fimbriatâ utrinque unidentatâ, ungue subcristato disco bidentato.

Sepala 1¾ *unciam longa, alba, sanguineo interruptè striata et fasciata.* Petala *æqualia et conformia, apice sanguinea basi maculata dorso alba.* Labellum *purpureum, ungue viridi cristâ dentibusque albis;* crista laciniæ intermediæ *fimbriata pilis submoniliformibus:* unguis *e cirrhulis* 4-6 *rectis submoniliformibus constans; dentes disci compressi divaricati margine crenati posteriore duplò majore.* Columna *basi viridis, apice lutea, et utrinque subalata, rostello longissimo subulato.* Caudicula *polliniorum longissima, glandulâ parvâ subtriangulari.*

For the opportunity of publishing this remarkably beautiful epiphyte, we are obliged to Mr. Knight, of the King's Road Nursery, with whom it flowered in the stove in July last.

It is a native of Trinidad, growing on old decayed branches of trees near the Mud Lake. We have seen a drawing of it taken on the spot, and it was in no degree superior in beauty to the present specimen.

The lip is a most curious organ. It is of a firm fleshy texture, and is deeply divided into three parts; of these the two lateral ones are falcate and smooth, while the middle one is rounded, has a strong tooth on each side, and is bordered by a deep white fringe of necklace-shaped hairs; then the short stalk of the lip has four or more such hairs growing straight from its border, and the disk is furnished with a large white two-lobed flat tooth, the divisions of which diverge from each other.

---

* See folio 1206.

The striping, banding, and painting of the delicate white flowers with rich crimson, produces a very rich and striking effect.

1812



S. Drake del.   Pub. by J. Ridgway 169 Piccadilly Nov. 1 1835.   J. Watts sc.

* GARDOQUÍA Gilliésii.

The Gillies Gardoquia.

DIDYNAMIA *GYMNOSPERMIA.*

Nat. ord. LABIACEÆ. (*Introduction to the Natural System of Botany, p.* 239.)

GARDOQUIA.—Suprà, vol. 21. *fol.* 1747.

G. *Gilliesii;* suffruticosa, ramis divaricatis pubescentibus, foliis oblongo-linearibus cuneatisve obtusis integerrimis basi angustatis utrinque viridibus planis, floralibus conformibus bracteisque calyce subbrevioribus, verticillastris paucis multifloris laxiusculis irregularibus, calycibus elongatis erectis, dentibus lanceolato-subulatis subæqualibus, fauce intùs nudâ. *Bentham Gen. & Sp. Labiat. p.* 403.
G. Gilliesii. *Graham in Edinb. Phil. Journ.* 1831, *Sept.*
G. chilensis. *Bentham in Hooker & Arn. Beech. Voy.* 58.

A neat little half-shrubby herbaceous plant, growing not more than 6 or 8 inches high, flowering from June to September in the open border, and readily multiplied by cuttings which root freely in peat and sand. It requires a little protection in winter.

It appears to be a common Chilian plant, for it occurs in almost every collection that has yet reached England from the neighbourhood of Valparaiso.

Our drawing was made in the Garden of the Horticultural Society.

Although this is far less showy than G. *Hookeri*, figured at fol. 1747, yet as it is a hardier plant it will be to many the more interesting of the two.

Our figure represents the analysis of the parts of fructification: 1. is a magnified entire flower; 2. is a calyx; 3. a corolla split open; and, 4. one of the anthers with a part of the filament.

---

* See folio 1747.

1813

1813.



[illegible] [illegible]

1813

* DAUBÉNYA aúrea.

Golden Daubenya.

HEXANDRIA *MONOGYNIA.*

Nat. ord. ASPHODELACEÆ. (*Introduction to the Natural System of Botany, p.* 273.)

DAUBENYA.—Inflorescentia umbellata, sessilis, epigæa. *Perianthium* tubulosum, ovario arcte appressum, limbo bilabiato, labio superiore nano tridentato, inferiore majore tripartito: in floribus radii maximo, disci depauperato. *Stamina* 6 inæqualia basi laciniarum adnata, declinata, in floribus disci sæpè submonadelpha. *Ovarium* elongatum, subtriangulare angulis rotundatis, in stylo angustatum; 3-loculare, loculis polyspermis apice vacuis. *Stigma* simplex.—— *Bulbi* capenses *habitu omnino* Massoniæ.

D. *aurea.*
Massonia lutea. *Hort.*

Folia 2, *oblonga, carnosa, sulcata, prostrata, umbellam sessilem multifloram e sinu suo emittentia.* Flores *lutei, densè aggregati;* radii *labio exteriore* (*inferiore*) *3-partito; laciniis obovatis rotundatis lateralibus brevioribus;* disci *laciniis omnibus angustis labii majoris acutis paulò elongatis.* Stamina 6, *basi laciniarum adnata, declinata, laciniæ inferiori intermediæ oppositum cum eâ altiùs connatum.* Antheræ *dorso affixæ.* Ovarium *laciniis nullis inter loculos.* Ovula *rotunda funiculo brevi placentæ adnata.*

A greenhouse bulb, native of the Cape of Good Hope, whence it was obtained by Messrs. Young of Epsom, under the name of *Massonia lutea.* It flowers in June, and is very pretty as well as singular, so long as its blossoms remain expanded, which is for about three weeks; afterwards its flaccid green leaves have little to attract attention.

No trace of this plant is to be found in books, so that it is probably some recent discovery made in the interior of the Cape Colony. From Massonia, with which it remarkably agrees in habit, it differs essentially in its tubular not campanulate, very irregular, perianth, and in the absence of the honey-pores which form so remarkable a part of the character of Massonia. With the fruit we are unacquainted.

---

* We name this new and curious genus after Dr Charles Daubeny, Professor of Botany at Oxford, whose interesting researches in Vegetable Chemistry have materially conduced to improve our knowledge of the physiology of plants.

Fig. 1. represents a flower of the ray with the larger lip; fig. 2. is one of the flowers of the disk, in which both lips are very small and imperfect; fig. 3. is a vertical section of the ovary, with the empty upper portion of the cells; fig. 4. is a transverse section of the ovary, shewing how closely it is invested by the tube of the perianth; this takes place to such a degree as to give the ovary the appearance of being inferior.

1814

1814.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Dec. 1. 1835. J. Watts sc.

* ELICHRÝSUM bícolor.

Two-coloured Elichrysum.

SYNGENESIA *POLYANDRIA.*

Nat. ord. ASTERACEÆ. (COMPOSITÆ.) *Introduction to the Natural System of Botany, p.* 197.*)*

ELICHRYSUM.—Suprà, vol. 1. *fol.* 21.

E. *bicolor;* annuum; foliis lineari-lanceolatis acuminatis basi obtusis scabro-ciliatis, superioribus subulatis, caule glabro ramoso ramulis monocephalis esquamatis, bracteis involucri fulvis aureisque acutis.

Caulis 2-*pedalis, sulcatus, fastigio ramoso.* Folia *basi aliquando subcordata, margine scabra, supra scabriuscula. Capitula facie omnia E. bracteati.*

A beautiful new hardy annual introduced by Mr. Low of the Clapton Nursery. In appearance it resembles the now common *E. bracteatum*, but is much handsomer.

It is a native of Van Diemen's Land, whence our excellent correspondent Mr. Gunn has sent beautiful specimens (No. 111), agreeing in all respects with the cultivated plant.

It is one of the prettiest new species of the season that has just past by.

---

* This is an old Greek name, meaning *a golden spiral,* and consequently should be written Helichrysum. It is supposed to have been the plant now called Gnaphalium Stæchas.

1815



Miss Drake del.   Pub by J. Ridgway 169 Piccadilly Dec 1. 1835.   J. Watts sc.

1815

[*] MACRADÉNIA triándra.

Triandrous Long-gland.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDACEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

MACRADENIA.—Suprà, vol. 8. *fol.* 612.

M. *triandra;* foliis coriaceis lineari-oblongis acuminatis, racemo prostrato, labello in medio trilamellato, clinandrio serrato antheris duabus sterilibus.

Pseudobulbi *cylindracei paulo attenuati.* Folia *plurivenia. Racemi foliis multò breviores.* Sepala *et* petala *intus sanguinea herbaceo limbata.* Labelli *lamellæ rotundatæ, costæformes, antrorsùm crassiores.* Antheræ steriles *ovales, integræ, sanguineo marginatæ.*

The *Macradenia lutescens* of this work (folio 612) we have never seen; but as it was described by Dr. Brown we have no difficulty in deciding, upon the evidence that the description affords, that it is specifically different from this.

No notice is taken by Dr. Brown of the existence of any lamellæ in the middle of the lip, nor of the abortive anthers which are always present; besides which the clinandrium here is regularly and strongly serrated, the leaves much narrower, and the raceme pendulous or prostrate instead of erect.

A native of Surinam, whence it was brought by Mr. Lance. It flowered in the possession of the Horticultural Society in May last. A strong damp stove heat is required to keep it in health.

---

[*] So named in allusion to the long caudicula of the pollen masses, fig. 5. from μακρος long, and αδην a gland.

1816

1816.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Dec 1 1835. J. Watts sc.

1816

* COCCOLÓBA virens.

Green Sea-side Grape.

OCTANDRIA *TRIGYNIA.*

Nat. ord. POLYGONACEÆ. (*Introduction to the Natural System of Botany*, p. 221.)

COCCOLOBA, L. *Calyx* 5-partitus, semi-inferus, laciniis imbricatis; in fructu incrassatus baccans. *Stamina* 7-10 serie simplici fauce inserta, filamentis basi in urceolum brevissimum coalitis. *Ovarium* semi-superum, trigynum; stigmatibus spathulatis nunc lobulatis. *Achenium* basi lobatum, osseum, calyce baccato tectum.——Frutices arboresve *minores, occidentales.* Flores *racemosi.* Fructus *subacidi, edules.*

C. *virens;* foliis ovato-lanceolatis obtusis basi in petiolum angustatis, racemis nutantibus, floribus decandris.

Folia *omninò depilata, ovato-lanceolata, semper basi angustata nec ullo modo basi obtusa vel cordata; omnia conformia.* Racemi *virides, nutantes, foliis minoribus æquales.*

The species of Coccoloba are so ill defined in books that it is by no means easy in the absence of fruit to determine them. This in some respects resembles C. *obtusifolia* of Jacquin, in others C. *microstachya* of Willdenow. From the former it differs in the form of its leaves, from the latter in their size and proportion to the racemes.

A hothouse plant, communicated from Wormleybury by Sir Abraham Hume, with whom it flowered in August 1833. It was called C. *excoriata*, which is a very different plant.

We are unacquainted with its native country, but presume it to be the West Indies.

---

* From κοκκος a fruit (in this case seed) and λοβος a lobe, in allusion to the lobed seeds.

1817



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Dec. 1. 1829. S. Watts sc.

* ÓXALIS Pióttæ.

Piotta's Oxalis.

DECANDRIA *PENTAGYNIA.*

Nat. ord. OXALIDACEÆ. (*Introduction to the Natural System of Botany, p.* 140.)

OXALIS.—Suprà, vol. 15. *fol.* 1249.

O. *Piottæ;* acaulis, petiolo planiusculo, foliolis 3 obcordatis glabris, scapo pubescente unifloro foliis longiore, sepalis integris, stylis longissimis, stigmatibus plumosis.
O. Piottæ. *Colla Hort. Rip. p.* 98. *t.* 1.

A truly beautiful little half hardy, or frame perennial, flowering most copiously during the months of July and August. A little tuft does not indeed produce much appearance, but a pot filled with its dense green leaves and covered with the large salmon coloured flowers is a lovely object.

Said to be a native of the Cape of Good Hope. Our specimens were obligingly communicated by Mrs. Marryat, to whom they were sent by Mrs. Palliser. That lady obtained them both from Professor Savi of Pisa and from the direction of other Italian Botanic Gardens.

We believe the plant is at present in the possession of no one in this country except Mrs. Marryat, and those of her friends to whom she has given it.

The species seems nearly allied to *O. compressa.*

---

* See folio 1249.

1818

1815.



Drake. del. Pub. by J. Ridgway 169 Piccadilly Dec. 1. 1835. J. Watts sc.

1818

* GALATÉLLA punctâta.

Dotted Galatella.

SYNGENESIA POLYGAMIA *FRUSTRANEA.*

Nat. ord. ASTERACEÆ. (COMPOSITÆ.) (*Introduction to the Natural System of Botany*, *p.* 197.)

GALATELLA, Cassini. *Capitulum* radiatum; radio uniseriali, ligulis fœmineis abortivis; disci flosculis hermaphroditis fertilibus tubulosis; stigmatibus apice spathulatis obtusis. *Involucrum* disco brevius; foliolis pluriserialibus imbricatis inappendiculatis; interioribus membranaceis exterioribus sæpè trinerviis. *Receptaculum* alveolatum, marginibus alveolorum paleaceo-laceris. *Pappus* simplex, pluriserialis, radiis filiformibus rigidis serrulato-scabris. *Achenia* densè hirsuta v. strigoso-setacea. *Nees Genera et Sp. Asterearum*, *p.* 158. (terminis mutatis.)

G. *punctata*; foliis lineari-lanceolatis acuminatis integerrimis trinervibus punctatis scabris, ramulorum linearibus, caule apice corymboso ramis elongatis paucifloris contractis, involucris brevibus turbinatis, radio longo. *Nees l. c.*
Aster punctatus. *W. & Kit. Pl. Hung.* 2. 113. *t.* 109.
Galatella intermedia. *Cass. in dict. des sc. nat. xviii. p.* 58.
Aster desertorum. *Fisch. ined.*

A hardy herbaceous plant, native of salt marshes in Hungary, Podolia, and elsewhere in the east of Europe. It also occurs in Siberia. In this country it flowers in July and August, growing to the height of about two feet, and forming a thick compact bush.

We greatly doubt whether it is really a distinct species from G. *hyssopifolia* and *acris*, or they from each other, so very difficult is it to distinguish them with absolute certainty when one has a long series of specimens under examination. From the former it may, however, be generally known by its compact inflorescence, and larger flower heads, which are quite regularly corymbose, instead of being of different heights as in *hyssopifolia;* its leaves too are more uniformly three-ribbed.

In *G. acris* the leaves are pretty uniformly one-ribbed, and the inflorescence much smaller and more contracted.

All the species of this genus are well adapted for borders of shrubberies, and for places where shade-loving plants alone will grow.

1819



Miss Drake del. Pub. by J. Ridgway 169 Piccadilly Dec. 1 1835. J. Watts sc.

1819

* OCHRÁNTHE arguta.

Fine-toothed Palebloom.

PENTANDRIA *TRIGYNIA.*

Nat. ord. HYPERICACEÆ ANOMALÆ.

OCHRANTHE. *Calyx* membranaceus, 5-phyllus, imbricatus, *corollæ* pentapetalæ simillimus. *Stamina* 5 hypogyna. *Discus* urceolatus pentagonus. *Carpella* 3, basi juncta, stylis disjunctis. *Ovula* cuique carpello 6, placentæ centrali affixa.——Folia *opposita, serrata.* Stipulæ *interpetiolares serratæ.* Flores *terminales, pallidi.*

Ochranthe arguta.

Caulis *fruticosus, glaber, adultus cinereus, junior viridis cinereo maculatus.* Folia *opposita, glabra, petiolata, obovato-lanceolata, acuminata, serrata, basi integra.* Stipulæ *intrapetiolares, ovatæ, serrulatæ, pallidæ.* Thyrsus *terminalis, congestus, pauciflorus.* Flores *inodori, albidi, demum flavescentes.* Pedicelli *basi* 1-, *medio* 2-*bracteati, glabri.* Calyx *erecto-patens, irregularis, 5-phyllus; sepalis concavis, obtusis, inæqualibus, exterioribus corolla brevioribus, margine subciliatis, æstivatione imbricatis* ⅖. Petala 5, *hypogyna, subconvoluta, rarò aliquot patentia, unguiculata, oblonga, obtusa, versus medium 3-venia.* Stamina 5, *hypogyna, erecta, rigida, pistilli longitudine, petalis alterna;* antheræ *medio affixæ, introrsæ, erectæ, biloculares, longitudinaliter dehiscentes;* pollen *rotundo-triquetrum, angulis (inflatis ?) globosis pellucidis, nunc subrotundum angulis nullis.* Discus *cyathiformis, pentagonus, carnosus, angulis planatis.* Ovarium *superum, ovatum, obtusè trigonum, stylis 3 subulatis erectis, versus basin pilis raris patentibus; 3-loculare, ovulis circiter 6 in utroque loculo, placentæ versus apicem axeos appensis.*

A native of China, and consequently a Greenhouse plant. It flowered in the Garden of the Horticultural Society so long since as March 1826; but shortly after died, and has never again made its appearance. We have kept the drawing ever since in our portfolio, in the hope of the species being again discovered, but we now find it necessary to publish some account of the plant.

---

* From ὠχρὸς pale, and ανϑος a flower.

The fruit being unknown, we are unable to obtain even an approximation to its true station in the system. In some respects it resembles Cunoniaceæ, especially in its remarkable stipulation, and apocarpous many-seeded pistil, combined with opposite leaves; but it differs in having only five hypogynous stamens, three instead of two carpels, and a remarkably imbricated calyx.

Eucryphia and Carpodontos among Hypericaceæ are polyandrous and have no stipules, besides other points of discrepancy.

Hugoniaceæ have alternate leaves, a different habit, and a larger number of stamens.

Anisadenia, which seems to be a shrubby Elatinaceous plant, has monadelphous stamens, and no hypogynous cup, &c. Its leaves are also alternate without stipules.

In all probability Ochranthe forms a part of some unknown order, the station of which will be in the Calycose group of polypetalous Dicotyledons, either in the Cistal or Guttal alliance. *(See Key to Structural, Physiological and Systematic Botany, p.* 51.*)*

1820

1820



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Dec. 1. 1835. J. Watts sc.

* RHODODÉNDRON pulchérrimum.

The lovely Rhododendron.

DECANDRIA *MONOGYNIA.*

Nat. ord. Ericaceæ. (*Introduction to the Natural System of Botany, p.* 182.)

RHODODENDRON.—Suprà, vol. 1. *fol.* 37.

Garden Variety.

A most beautiful plant obtained by Mr. Waterer of Knaphill between R. arboreum and caucasicum. It is of rather delicate appearance, but we are informed that it is quite hardy, and an abundant flowerer. Our drawing was made in March last.

Fig. 1 is a kind called Rhododendron Nobleanum, which is very much like the other in all respects, except that its flowers are of a deep and brilliant rose colour.

Both are among the handsomest hardy shrubs in cultivation.

---

* See folio 1240.

1821

1821.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Dec. 1. 1835. S. Watts sc.

1821

* EULÓPHIA lúrida.

Lurid Eulophia.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDACEÆ, § VANDEÆ, *Lindl.* (*Introduction to the Natural System of Botany, p.* 262.)

EULOPHIA.—Suprà, vol. 17. *fol.* 1433.

E. *lurida;* foliis (lineari-) lanceolatis scapo ramoso multò brevioribus, bracteis minimis subulatis, sepalis lineari-spatulatis obtusis, petalis paulò latioribus, labelli tripartiti basi callosi lobis lateralibus obtusis recurvis: intermedio obcordato, calcare cylindraceo inflexo obtuso. *Lindl. gen. et sp. Orch. p.* 182.

One of the easiest of all stove Orchideous plants to cultivate, growing upon a damp wall, or a rough stone, or under almost any kind of condition, provided the air be hot and damp, and uniform. It then flowers profusely at intervals throughout the whole year. Our drawing was made in the noble collection of the Messrs. Loddiges in January last.

A native of Sierra Leone, whence great quantities are occasionally brought. It appears to be extremely common upon the trunks of trees in some parts of that colony.

---

* See folio 1433.

1822



Miss Drake. del. Pub by J. Ridgway 169 Piccadilly Jan. 1. 1836. J. Watts sc.

1822

* COSMÉLIA rúbra.

Red Cosmelia.

PENTANDRIA *MONOGYNIA.*

Nat. ord. EPACRIDACEÆ.

COSMELIA. *Calyx* foliaceus. *Corolla* tubulosa. *Stamina* epipetala; *antheræ* apicibus ciliatis filamentorum adnatæ. *Squamulæ* 5, hypogynæ. *Capsula* placentis columnæ centrali adnatis.——Frutex *paludosus, erectus, ramosus,* ramis *dum denudatis non cicatrisatis.* Folia *basibus cucullatis semivaginantibus.* Flores *ramos laterales breves terminantes, solitarii, cernui.* Calyx *foliis minoribus imbricatus.* Corolla *ruberrima.* Stamina *inclusa, antheris basi solutis.* Placentæ *utraque extremitate liberæ.* Brown prodr. 553.

C. rubra. *Brown l. c.*

According to Dr. Brown this is a marsh plant, found upon the South Coast of New Holland, and remarkable for its leaves, although so thick and rigid, not leaving conspicuous scars upon the branches when they fall off.

In this country it is a very pretty greenhouse plant with the habit of an Epacris. Our drawing was made in Messrs. Loddiges collection in May last.

The dissected parts of the flower in the accompanying plate afford a good illustration of the Natural order Epacridaceæ, which are only known from true Ericaceæ by their anthers having single cells opening by a pore. 1. Is the top of a corolla cut open, with the stamens sticking to it; 2. an anther with a portion of the upper end of a filament; 3. a pistil, with the five hypogynous scales.

---

* Apparently from *κοσμεω* to ornament, in allusion to its beauty.

1823

1823.



Miss Drake del. Pub by J. Ridgway 169 Piccadilly Jan. 1. 1836. J. Watts sc.

* LASTHENIA califórnica.

Downy Lasthenia.

SYNGENESIA POLYGAMIA *SUPERFLUA.*

Nat. ord. ASTERACEÆ (COMPOSITÆ).
LASTHENIA.—Suprà, vol. 21. *fol.* 1780.

L. *californica;* subpubescens; foliis integerrimis, capitulis basi umbonatis. *Supra l. c.*
L. californica. *De Cand. prodr. ined.*

This species differs from the one already published in the present work in little except its smaller size, and differently formed flower heads, which are impressed at the base with a cavity for the peduncle, and like the peduncles themselves are slightly downy.

It is an annual plant, flowering for about six weeks at different periods of the year, according to the season at which its seeds are sown.

Professor De Candolle, in a manuscript list of the genera of Compositæ, which we have just received from him, includes it in his first series *Tubulifloræ*, fourth tribe *Senecionideæ*, fifth sub-tribe *Heleniеæ*, first division *Gaillardieæ*, second sub-division *Euhelenieæ*, along with Cephalophora and Helenium, both of which are known in our gardens. We observe that Professor De Candolle attributes the genus to Cassini, but we have not succeeded in discovering it in any of the writings of that most unmethodical of Botanists.

---

* See folio 1780.

1824

1824



Miss Drake. del.

Pub. by J. Ridgway 169 Piccadilly Jan. 1. 1836.

J. Watts

* ARISTOLOCHÍA fœtens.

Stinking Birthwort.

GYNANDRIA *HEXANDRIA.*

Nat. ord. Aristolochiaceæ.
ARISTOLOCHIA.—Suprà, vol. 8. *fol.* 689.

A. *fœtens;* foliis lato-cordatis acutis, caule volubili, pedunculis solitariis, bracteâ perfoliatâ, limbo calycis maximo integro cordato labio longissimè caudato, tubo extus glabro.

Nauseosa, fœtida, demum semiputrida ramis scandit longis volubilibus glabris. Folia *cordata sinu aperto, integerrima, subrotunda, acuminata, subtus levissimè pubescentia, glaucescentia.* Bracteæ *solitariæ, orbiculatæ, perfoliatæ.* Flores *maximi, tubo extus glabro, basi inflato, sursum angustato costato arcuato, limbo subrotundo cordato basi clauso apice in appendicem longum linearem subspiralem acuminatum producto, luteo purpureoque livido variegato maculato et punctato.*

A native of the West Indies, whence it has been obtained by Mrs. Marryat, in whose stove at Wimbledon it flowered in June last. Like all the other hothouse species it strikes readily from cuttings.

It is chiefly remarkable for the large size, and singular colour of its flowers, which are beautifully variegated with purple and dirty yellow; they have a most disagreeable disgusting smell, which will prevent the plant from becoming a favourite.

Nearly allied to A. *grandiflora,* from which it is distinguished by the smooth tube of its calyx, the colour of the same part, &c. &c.

---

* See folio 1399.

1825





Miss Drake del. Pub by J. Ridgway 169 Piccadilly Jan 1 1836 J. Watts sc.

* PLEUROTHÁLLIS pícta.

Painted Pleurothallis.

GYNANDRIA *MONANDRIA.*

Nat. ord. ORCHIDACEÆ, § MALAXIDEÆ.
PLEUROTHALLIS.—Suprà, vol. 9. *fol.* 759.

P. *picta;* folio spathulato marginato retuso racemis laxis duplò breviore, bracteis minimis, sepalis acuminatis lateralibus apice tantùm sejunctis, petalis lineari-lanceolatis acutis, labello lineari obtuso carnoso supra 1-sulcato. *Supra fol.* 1797.

Very near the *P. Grobyi* already figured in this work (fol. 1797), from which it principally differs in its spathula-shaped retuse leaves, and in the narrower form of all the parts of the flower; as will be seen by comparing the dissectiòns in the two plates.

This also is a native of Demerara, and was sent us by Mr. Loddiges first in October 1834, and next in March 1835, when our drawing was made. It is a graceful pretty species, and well deserves an attentive examination.

Its tufts are rapidly formed under good management, but it seems to require the close atmosphere of a bell-glass.

Fig. 1. is a view of the column, petals, and lip. 2. The column by itself. 3. The upper side of the lip. 4. The pollen mass in situ. 5. The same separated from each other.

---

* See folio 1298.

1826